

Journal Pre-proof

Efficacy and Safety of Pegylated Interferon alfa-2b in Moderate COVID-19: A phase II, randomized, controlled, open-label study

Anuja Pandit, Nirav Bhalani, B.L. Shashi Bhushan, Parshottam Koradia, Shweta Gargiya, Vinay Bhomia, Kevinkumar Kansagra



PII: S1201-9712(21)00232-0
DOI: <https://doi.org/10.1016/j.ijid.2021.03.015>
Reference: IJID 5217

To appear in: *International Journal of Infectious Diseases*

Received Date: 12 January 2021
Revised Date: 2 March 2021
Accepted Date: 5 March 2021

Please cite this article as: { doi: <https://doi.org/>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Efficacy and Safety of Pegylated Interferon alfa-2b in Moderate COVID-19: A phase II, randomized, controlled, open-label study

Authors: Anuja Pandit, MD^a, Nirav Bhalani, MD^b, B L Shashi Bhushan, MD^c, Parshottam Koradia, MD^d, Shweta Gargiya, MBBS^e, Vinay Bhomia, MD^f, Kevinkumar Kansagra, MD^g

Affiliations: ^a National Cancer Institute, Badsa, Jhajjar, Haryana 124105, India; ^b Rhythm Heart Institute, Near Siddharth Bungalows, Sama Savli Road, Vadodara 390022, Gujarat, India; ^c Victoria Hospital, Bangalore Medical College & Research Institute, Fort K R Road, Bangalore Urban, Karnataka 560002, India; ^d BAPS Pramukh Swami Hospital, Shri Pramukh Swami Maharaj Marg, Adajan Char Road, Adajan, Surat 395009, Gujarat, India; ^e Tapan Research Centre, Basement Tapan Hospital Nr. Platinum Hall, Anandnagar Cross Road Satellite, Ahmedabad 380015, India; ^f Sanjivani Super Speciality Hospital, 1, Uday Park Society, Nr. Sunrise Park, Vastrapur, Ahmedabad 380015, Gujarat, India; ^g Zydus Research Centre, Clinical R & D, Cadila Healthcare Limited, Sarkhej-Bavla N. H. No. 8 A, Moraiya, Ahmedabad-382213, Gujarat, India

Corresponding author:

Dr. Kevinkumar Kansagra, MD

Zydus Research Centre, Clinical R & D, Cadila Healthcare Limited,

Sarkhej-Bavla N. H. No. 8 A, Moraiya, Ahmedabad-382213, Gujarat, India

Tel: +91-2717-665555, Ext.: 279

Fax: +91-2717-665355

Mobile: +91-9898123647

E-mail: kevinkumarkansagra@zyduscadila.com

HIGHLIGHTS

- Type I interferons-a/b exhibiting direct inhibitory effects on viral replication.
- We evaluated efficacy and safety of PEG IFN- α 2b in moderate COVID-19 subjects.
- Study provides initial evidence for the potential use of PEG IFN- α 2b in COVID-19.
- PEG IFN- α 2b was shown to significantly improve the clinical outcome.
- PEG IFN- α 2b reduces duration of viral shedding as per Phase 2 data.

ABSTRACT

Objective: To evaluate the efficacy and safety of pegylated interferon alfa-2b (PEG IFN- α 2b) along with the standard of care (SOC) in subjects with moderate COVID-19.

Methods: In this phase 2, randomized, open-label study, adult subjects aged >18 years with RT-PCR confirmed COVID-19 with moderate symptoms were randomized in a 1:1 to receive PEG IFN- α 2b plus SOC, or SOC alone. The primary endpoint was improvement in clinical status on day 15, measured by WHO 7-point ordinal scale.

Results: Total 40 subjects were randomized to PEG IFN- α 2b plus SOC (n=20) and SOC (n=20). Overall, 19 (95.00%) subjects in PEG IFN- α 2b plus SOC had achieved clinical improvement on day 15 compared to 13 (68.42%) subjects in SOC ($p < 0.05$). Overall, 80% and 95% of subjects in the PEG IFN- α 2b plus SOC group had a negative RT-PCR result on day 7 and day 14, respectively as compared to 63% and 68% in the SOC group. Adverse events (AEs) were reported for 11 subjects in the PEG IFN- α 2b plus SOC group and 8 subjects in the SOC group. All reported AEs were mild.

Conclusion: The significant improvement in clinical status on day 15 is likely due to faster viral reduction compared to SOC with the PEG IFN- α 2b treated moderate COVID-19 subjects showing a difference as early as day 7 and becoming significant by day 14.

Keywords: Antiviral, coronavirus, pegylated interferon alfa-2b (PEG IFN- α 2b), COVID-19

Introduction

A novel coronavirus disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged in December 2019 (COVID-19) in a cluster of patients in Wuhan, China, which has been designated a worldwide pandemic (Cucinotta and Vanelli, 2020; Spinelli and Pellino, 2020). As of 31 January 2021, there have been 102,139,771 confirmed cases of COVID-19 worldwide, including 2,211,762 reported deaths (WHO, 2021).

Bats are the zoonotic reservoir of several coronavirus (CoV) strains and several viruses. Bats and viruses have been co-existing and co-evolving over millions of years. The sequence of SARS-CoV-2 is similar to bat severe acute respiratory syndrome (SARS)-like CoV. One of the effects of a strong evolved immune mechanism or molecules of the primary innate immunity is the interferon. Interferons play a significant role in the controlling mechanism of viral replication. In bats, different types of interferons such as Type I or Type II interferons have been recognized. The evolved immune mechanisms help them to harbor viruses without any clinical symptoms (Chakraborty et al., 2020). Most of the patients with COVID-19 develop seroconversion between 7 and 14 days after diagnosis (Vabret et al., 2020; Zhao et al., 2019). COVID-19 is effectively transmitted from human to human, with influenza-like symptoms ranging from mild disease to severe disease and multi-organ failure, eventually resulting in death, especially in aged patients (≥ 50 years) with co-morbid conditions (Zhou et al., 2020; Zhang J et al., 2020).

Type I interferons- α/β are broad spectrum antivirals, exhibiting both direct inhibitory effects on viral replication and supporting an immune response to clear virus infection (Wang and Fish, 2019). Pegylated interferon alfa-2b (PEG IFN- α 2b) is a covalent conjugate of recombinant α 2b interferon with monomethoxy polyethylene glycol. It binds to and activates human type 1 interferon receptors causing them to dimerize. This activates the JAK/STAT pathway. Activation of the JAK/STAT pathway increases expression of multiple genes in

multiple tissues involved in the innate antiviral response. There are published results reporting the role of interferon in the treatment of SARS-CoV and middle east respiratory syndrome coronavirus (MERS-CoV) (Perlman and Dandekar, 2005; Stroher et al., 2004; Falzarano et al., 2013). Interferon- α and its pegylated form have been used clinically in the treatment of Hepatitis B and C viruses for a number of years. Interferons inhibit viral infection by inducing both innate and adaptive immune responses like altering the intracellular environment to restrict viral replication, and inducing signaling events that activate immune cell populations and thereby elicit an antiviral immune response (Loutfy et al., 2003).

Literature demonstrates that the dynamics of interferon -related antiviral responses could lower the virulence of the current COVID-19 outbreak (Nezhad et al., 2020). Perhaps the most exciting support for the potential benefit of interferon alfa in COVID-19 comes from the publication of two research articles. The first one by Lokugamage et al exhibited a direct anti-viral effect of interferon alfa against the novel coronavirus in vitro. The study demonstrated around 10,000 fold reduction in virus titre in cells that were pre-treated with interferon alfa 48 hours earlier (Lokugamage et al., 2020). The second by Zhou et al retrospectively analyzed 77 moderate COVID-19 subjects in Wuhan and observed that those who received interferon- α 2b showed a significant reduction in the duration of the virus shedding period (accelerated viral clearance by \sim 7 days from onset of symptoms) which was co-related with reduced levels of inflammatory cytokine, IL-6 (Zhou Q et al., 2020). This suggests that providing PEG IFN- α 2b to COVID-19 patients earlier in the disease is expected to make the patients viral-free sooner and reduce their chances of deteriorating to severe states of disease.

We performed a multi-centric, randomized, open-label study to evaluate the efficacy and safety of a single dose of PEG IFN- α 2b in addition to standard of care (SOC) compared to SOC alone in patients with moderate COVID-19.

Materials and Methods

Study Design

This phase 2, multi-centric, randomized, open-label study evaluated the efficacy and safety of a single dose of PEG IFN- α 2b in the treatment of adult subjects diagnosed with SARS-CoV-2, and the study was undertaken at 6 study centers in India. Eligible subjects were randomly assigned in a 1:1 ratio to receive either PEG IFN- α 2b along with SOC, or SOC alone.

This study was initiated after obtaining the approvals of the Ethics Committee (EC) at each site, Drugs Controller General of India (DCGI) (dated 08 May 2020) and was overseen by an Independent Data Safety Monitoring Board. This study was conducted in accordance with the applicable local regulations and registered with the CTRI (CTRI identifier: CTRI/2020/06/026087).

Study Populations

Individuals with suspected COVID-19 were recruited from 6 study centers across India from 08 July 2020 to 04 September 2020. Key inclusion criteria were age ≥ 18 years, RT-PCR confirmed SARS-CoV-2 infection, pneumonia with no signs of severe disease, respiratory rate 15 to 30 breaths/minute, SpO₂ 90% to 94%, and for female patients of child-bearing potential, a negative pregnancy test prior to treatment. Additional inclusion criteria included C-reactive protein (CRP) < 16 mg/L, IL-6 < 100 pg/mL, D-dimer < 2 μ g/mL, interferon- γ , ferritin, tumor necrosis factor (TNF)- α , IL-1 β greater than upper limit of normal (ULN), illness of any duration and radiographic infiltrates by chest x-ray or evidence of rales/crackles or other clinical symptoms on clinical examination.

Key exclusion criteria were alanine aminotransferase (ALT)/aspartate aminotransferase (AST) $>5 \times$ ULN, stage 4 severe chronic kidney disease or required dialysis (i.e. estimated glomerular filtration rate <30 mL/min/1.73 m²), pregnant or breast-feeding women, severe co-morbidity (e.g. uncontrolled hypertension, uncontrolled diabetes mellitus, systemic disease which had affected the vital organs severely, immunocompromised patients etc.), comorbid condition like myocardial infarction or heart failure within 90 days of recruitment, and prolonged QT interval (>450 ms).

Interventions

Eligible subjects were randomized in a 1:1 ratio to either PEG IFN- α 2b (1 μ g/kg subcutaneous [SC] injection, single dose) plus SOC, or SOC alone. The regulatory recommendations (Clinical Management Protocol: COVID-19 [Ministry of Health and Family Welfare, 2020]) have been followed to categorize moderate COVID-19 subjects and treatment accordingly. During the study, all the investigators were in agreement to provide standard of care to all the subjects. Antipyretics, cough suppressants, antibiotics, steroids, vitamins, anticoagulant and hydroxychloroquine were administered as per regulatory recommendation and approval. Randomization was generated using SAS[®] software (Version 9.4). All subjects were hospitalized, RT-PCR tests in pharyngeal swab performed on screening, on day 7 and on day 14 and were discharged only after 2 consecutive negative RT-PCR tests and clinical cure.

Assessments

The primary efficacy endpoint was clinical status assessed on day 15 on a WHO 7-point ordinal scale consisting of the following categories: 1, not hospitalized, no limitations of activities; 2, not hospitalized, limitation on activities; 3, hospitalized, not requiring supplemental oxygen; 4, hospitalized, requiring supplemental oxygen; 5, hospitalized, on

non-invasive ventilation or high flow oxygen devices; 6, hospitalized, on invasive mechanical ventilation or extracorporeal membrane oxygenation (ECMO) ; and 7, death.

The secondary efficacy endpoints were the proportion of subjects with adverse events (AEs) that occurred on or after the first dose of PEG IFN- α 2b for up to 29 days, qualitative PCR for SARS-CoV-2 in pharyngeal swab, occurrence and duration of supplemental oxygen and mechanical ventilation, duration of hospitalisation, change from baseline in white blood cell count (WBC), haemoglobin (Hb), platelets, creatinine, glucose, total bilirubin, ALT and AST, and change from baseline in CRP, IL-6, D-dimer, interferon- γ , ferritin, TNF- α , and IL-1 β till Day 14.

Safety assessments were based on physical examinations, vitals, laboratory tests and the incidence and severity of AEs.

STATISTICAL ANALYSIS

The study was initiated in rapid response to the COVID-19 public health emergency, at which time there was very limited information about clinical outcomes in subjects with COVID-19.

There was no formal calculation of sample size for this study. A total of 40 subjects were enrolled in the study; 20 subjects each in PEG IFN- α 2b plus SOC and SOC alone.

The primary efficacy endpoint was the proportion of subjects showing improvement in condition (clinical status) measured using the WHO 7-point ordinal scale for clinical improvement during the dosing period, and was presented descriptively as frequency and percentage. Improvement is defined as a score of less than 2 on the WHO 7-point ordinal scale. Treatment effect was assessed using Fisher's exact test for active treatment (PEG IFN- α 2b plus SOC) versus SOC. Non-parametric test Wilcoxon rank sum test was used to assess the change in score from baseline within the group.

Secondary endpoints, qualitative RT-PCR, requirement/duration of supplemental oxygen, and mechanical ventilation were analyzed the same way as the primary endpoint by non-parametric Wilcoxon Rank Sum Test. Comparison of the following secondary endpoints, laboratory parameters and biomarkers (CRP, IL-6, D-dimer, interferon- γ , ferritin, TNF- α , and IL 1- β) between treatment groups were analyzed using an ANCOVA model treatment as fixed effect and baseline value as covariate.

Statistical significance was tested at a two-sided p-value of 0.05 unadjusted for multiple comparisons. Results are presented as mean \pm SD (in the text and tables).

Efficacy analyses were performed according to the modified intent-to-treat (mITT) population and Per Protocol (PP) population supportive for primary endpoint. The mITT population included all randomized subjects who received either of the study medication and appeared for at least one post-baseline efficacy assessment. The PP population included all randomized subjects who met the eligibility criteria, completed study in compliance with the protocol and did not have major protocol deviations. Safety analyses were performed using safety population, defined as all randomized subjects who received at least one dose of the study medication.

Results

Subject Disposition and Characteristics

A total of 86 subjects were screened and 40 subjects were randomized in the study. Out of 40, 39 subjects completed the study (20 subjects in the PEG IFN- α 2b plus SOC and 19 subjects in the SOC group). One subject in the SOC group discontinued from the study due to withdrawal of consent.

Of the 40 subjects randomized, 39 (97.50%) subjects comprised the mITT and PP populations, respectively.

Of the 40 subjects, 30 (75.00%) were male and 10 (25.00%) were female. The mean age was 49.35 ± 14.89 years in the PEG IFN- α 2b plus SOC group and 49.10 ± 12.44 years in the SOC group alone. Overall, demographic characteristics of the study subjects were comparable across the treatment groups (Table 1). The subject disposition is provided in Figure 1.

Primary Endpoint

The primary outcome (status on the WHO 7-point ordinal scale on day 15) was assessed in all subjects who were still in the hospital on day 15 exactly, and in outpatients (by means of telephonic follow-up) as close to day 15 as possible.

In both the populations, 19 (95.00%) and 13 (68.42%) subjects had achieved clinical improvement in PEG IFN- α 2b plus SOC and SOC group alone respectively on day 15 (Table 2). There was a statistically significant difference observed in clinical improvement in the PEG IFN- α 2b plus SOC group compared to the SOC alone from day 0 to day 15 ($p < 0.05$).

Subjects in the PEG IFN- α 2b plus SOC group achieved a higher reduction in mean score (measured by WHO 7-point ordinal scale) from baseline to day 15 than the subjects in the SOC group. The mean (SD) change in score from baseline to day 15 was $-2.25 (0.55)$ and $-2.05 (0.85)$ in the PEG IFN- α 2b plus SOC group and the SOC group, respectively (Table 3).

Secondary Endpoints

Of the 20 subjects, 16 (80.00%) and 19 (95.00%) subjects were tested negative in the RT-PCR in the PEG IFN- α 2b plus SOC group on day 7 and day 14 respectively (Figure 2). While in the SOC group alone, out of 19, 12 (63.16%) and 13 (68.42%) subjects were tested negative in the RT-PCR on day 7 and day 14 respectively. There was a significant statistical difference observed on day 14 between the PEG IFN- α 2b plus SOC and the SOC groups ($p < 0.05$).

Subjects in the PEG IFN- α 2b plus SOC group had a shorter duration of supplemental oxygen than subjects in the SOC group alone (Figure 3), median, 33.96 hours in the PEG IFN- α 2b plus SOC group, as compared to 49.75 hours in the SOC group; $P>0.05$. None of the subjects required mechanical ventilation during the study. Subjects were observed to have a similar duration of hospitalization in both the treatment groups during the study (median: 8 days, $P>0.05$).

Serial laboratory measurements of blood levels for WBC, Hb, platelets, creatinine, glucose, total bilirubin, ALT, AST, CRP, IL-6, D-dimer, interferon- γ , ferritin, TNF- α and IL 1- β were also conducted. There were no significant differences observed between the treatment groups for any of these parameters during the study (See Supplementary Table 1 to Table 8).

Safety

A total of 19 subjects reported at least one AE during the study: 11 subjects in the PEG IFN- α 2b plus SOC group and 8 subjects in the SOC group alone (see Table 4). All AEs were mild in severity. None of the subjects were discontinued from the study due to AEs in any of the treatment groups. There were no deaths and serious adverse events (SAEs) reported during the study period. All AEs were followed up until the subject 'recovered' or 'recovered with sequelae' or until the end of post treatment follow-up, whichever came first.

The most frequently reported AEs (Table 4) in the PEG IFN- α 2b plus SOC group were headache: 8 (40.0%), vomiting: 6 (30.0%), difficulty in breathing: 2 (10.0%), breathlessness: 1 (5.0%), dryness in mouth: 1 (5.0%), hypoxia: 1 (5.0%), and nausea: 1 (5.0%). The most frequently reported AEs in the SOC group alone were headache: 4 (20.0%), breathlessness: 3 (15.0%), chest pain: 2 (10.0%), difficulty in breathing: 2 (10.0%), dry mouth: 2 (10.0%), vomiting: 2 (10.0%), dryness in mouth: 1 (5.0%) and nausea: 1 (5.0%).

No apparent difference was observed in any of the lab parameters between the treatment groups. No clinically relevant findings from clinical examination, vital signs and ECG evaluations were attributed to PEG IFN- α 2b. Overall, single dose of PEG IFN- α 2b was safe and well-tolerated in the study.

Discussion

Recently (Hadjadj et al., 2020), an integrated immune analysis carried out on a cohort of 50 COVID-19 patients with various levels of disease severity revealed that the severe and critical patients were associated with a phenotype characterized by a highly impaired interferon type I response (associated with no interferon- β expression and low interferon- α production and activity), a persistent blood viral load and an exacerbated inflammatory response suggesting that the impaired type I interferon activity may be responsible for severe disease in COVID-19 patients. The important role of type I interferons such as interferon- α 2b in the disease severity of COVID-19 has been further illustrated by two additional reports, one (Bastard et al., 2020) showing that about 10.2% of patients with life threatening COVID-19 disease had neutralizing antibodies against type I interferons making them ineffective, and the other (Zhang Q et al., 2020) showing that around 3.5% patients with life threatening COVID-19 pneumonia had genetically defective induction and amplification of type I interferons. The relatively higher disease severity observed in SARS-CoV-2 infection in comparison to that observed in other respiratory infections may also be due to the relatively lower levels of induction of type I and type III interferon responses induced by the former (Blanco-Melo et al., 2020). Given that expression of type I interferons early in the infection helps not only in reducing both viral replication and secondary viral infection of neighboring cells but also in the activation and development of innate and adaptive antiviral immunity, an early intervention of COVID-19 patients with recombinant interferon- α 2b appeared to provide a realistic possible treatment in the management of this disease. Such an intervention

would not only be expected to reduce the overall viral burden but also reduce infection-related tissue damage.

In COVID-19, duration of viral shedding and viral load kinetics are important determinants for early disease transmission. The mean shedding time of RNA was 17.0 days (95% CI 15.5 - 18.6; 43 studies, 3229 individuals) in the upper respiratory tract. The drug with anti-viral property will help to reduce the duration of viral shedding and as well as reduce the kinetics of viral load (Cevik et al., 2020).

Treatment with interferon α 2b with or without arbidol was reported to significantly reduce the duration of detectable virus in the upper respiratory tract and the time period of elevated levels of the inflammatory markers IL-6 and CRP in blood (Zhou et al., 2020). This study further strengthened the idea of testing interferon α 2b in treating COVID-19. Historically, the pegylated form of interferon- α 2b because of its extended half-life in the body has been demonstrated to have a significantly higher efficacy in comparison to standard non-pegylated interferon alfa in the treatment of chronic Hepatitis C (Lindsay et al., 2001; Poynard et al., 2002). In Hepatitis C patients, treatment-induced improvement in liver necrosis and inflammation ranged from 39% (interferon) to 73% (pegylated interferon and ribavirin; $P < 0.001$). Significant reduction in worsening of fibrosis was also observed - 23% (interferon) and 8% (pegylated interferon and ribavirin; $P < 0.001$) (Poynard et al., 2002). Similarly, all three tested PEG IFN- α 2b doses (0.5, 1.0, or 1.5 $\mu\text{g}/\text{kg}$) significantly ($P \leq .042$) improved virologic response rates (i.e., loss of detectable serum hepatitis C virus [HCV] RNA) after treatment and after follow-up, as compared to interferon- α 2b (Lindsay et al., 2001). This knowledge led us to test the PEG IFN- α 2b product instead of the standard non-PEG IFN- α 2b in the treatment of moderate COVID-19 patients.

This randomized-controlled open-label study showed that PEG IFN- α 2b along with SOC may be beneficial in the treatment of subjects with moderate COVID-19. A single dose of PEG

IFN- α 2b administered along with SOC was shown to significantly improve the clinical status in moderate COVID-19 subjects when compared to those given SOC alone. Overall, 95% subjects in the PEG IFN- α 2b group showed an improvement of clinical status on day 15 as compared to only 68% subjects in the SOC.

This study further demonstrated that the PEG IFN- α 2b induced viral clearance was also faster than that achieved with SOC only. Overall, 80% and 95% of subjects in the PEG IFN- α 2b group had a negative RT-PCR result on day 7 and day 14 respectively as compared to 63% and 68% subjects in the SOC group. There was a statistically significant difference observed in qualitative RT-PCR result on day 14 between the treatment groups. The faster viral clearance may reduce the duration of infectivity of these patients, further resulting in a reduced secondary attack rate and shorter quarantine period, and also reduce community transmission (Schiffer et al., 2020).

A few subjects in both the treatment groups of the study required oxygen support during the course of their treatment. However, subjects in the PEG IFN- α 2b group needed supplemental oxygen for a shorter duration than those in the SOC group. None of the subjects required mechanical ventilation in either of the two treatment groups in the study. Also, subjects in the PEG IFN- α 2b group had comparable median duration of hospital stay compared with those in the SOC group. However, this small study clearly showed that treatment with PEG IFN- α 2b may have prevented disease progression to severe respiratory disease, and averted respiratory disease related complications.

Overall, PEG IFN- α 2b given along with SOC was found to be safe and well-tolerated in this study. There were neither any serious adverse events nor adverse events that led to drug discontinuation or death. All adverse events observed in the study were mild in severity. The most commonly reported treatment-related adverse events during clinical trials with PEG IFN- α 2b in combination with ribavirin in adults patients with HCV were fatigue, headache,

injection site reaction, nausea, chills, insomnia, anaemia, pyrexia, myalgia, asthenia, pain, alopecia, anorexia, weight decreased, depression, rash and irritability. Fatigue, alopecia, pruritus, nausea, anorexia, weight decrease, irritability and insomnia occurred at a notably lower rate in patients treated with PEG IFN- α 2b monotherapy when compared to those treated with combination therapy. In our study, the reported AEs in the PEG IFN- α 2b plus SOC treatment group in patients with COVID-19 were headache, vomiting, breathlessness, dryness in mouth, hypoxia and nausea. Therefore, the reported adverse events in this study are in line with the safety profile of PEG IFN- α 2b used for its approved indications.

The improvement in clinical status of moderate COVID-19 subjects reported in this study is also consistent with that of some other anti-viral drugs published in literature. Spinner et al., 2020 conducted a randomized controlled study to evaluate the clinical benefit of 5-day and 10-day courses of remdesivir compared to SOC in subjects with moderate COVID-19. On day 14, the 5-day and 10-day remdesivir groups showed significant improvement in clinical status as compared to the SOC group. A total of 76% of subjects in the 5-day remdesivir group and 77% of subjects in the 10-day remdesivir group had achieved clinical improvement on day 14, as compared to 68% of subjects in the SOC group (Spinner et al., 2020). While our observations are similar to those reported in the above study, the latter come from a much larger sample size.

Thus, our study indeed has some limitations which have been discussed here. First, the study cohort was small with a total of 40 subjects (20 subjects in each group). Second, subjects were followed-up for only 29 days after their first dose. In this regard, a Phase 3 study using PEG IFN- α 2b in the treatment of moderate COVID-19 patients is already in progress in India where the sample size is 250 subjects and the follow-up duration is 29 days. Another limitation of the current Phase 2 study may be that we did not analyze the pharmacokinetic (PK) profile of the drug and did not test its immunogenicity. In this regard, the PEG IFN- α 2b

used in the current study had already been tested in healthy volunteers in a Phase 1 setting where this biosimilar drug was compared to the innovator drug PegINTRON[®] in terms of both PK and immunogenicity, and found to be comparable. Further, a single dose administration of PEG IFN- α 2b may not be expected to cause immunogenicity concerns. However, the ongoing Phase 3 study would certainly address the safety and efficacy parameters more deeply. Subjects with comorbid conditions were not randomized as well as the cohort consisted of only moderate cases of COVID-19. This study was conducted to establish the proof of concept (POC) of using PEG IFN- α 2b in COVID-19 as an antiviral agent to improve clinical score at the tested dose (1 μ g/kg) while not causing any cytokine storm. While this POC was clearly established in the current study, the already ongoing Phase 3 clinical study in India will evaluate the safety and efficacy of PEG IFN- α 2b in a much larger sample size supporting an eventual market authorization of the drug in India.

Conclusion: This study provides initial evidence for the potential use of a single 1 μ g/kg dose of PEG IFN- α 2b in the treatment of moderate COVID-19 disease. The significant improvement in clinical status on day 15 is likely due to faster viral reduction compared to SOC with the PEG IFN- α 2b treated subjects showing a difference as early as day 7 and becoming significant by day 14. The absence of AEs in moderate COVID-19 patients suggests that this anti-viral drug may even be tested in early-disease patients. Treatment with PEG IFN- α 2b may also benefit in slowing the tide of this pandemic by reducing the duration of viral shedding. Further confirmatory studies are required to support the evidences observed in this study.

Author Contributions

Deven Parmar, Kevinkumar Kansagra, and Sanjeev Kumar Mendiratta were involved in conceptualization of the study. Hemal Mistry and Jatin Patel were involved in data

interpretation, manuscript writing, and manuscript review. Sunil Sharma was involved in statistical analysis, designing, programming and generation of TLFs and aided in interpretation of results. Purav Trivedi, Abhijit Mali, Brilina Patel, Lokesh Bathula and Vishal Nakrani provided operational support. Anuja Pandit, Nirav Bhalani, B L Shashi Bhushan, Parshottam Koradia, Shweta Gargiya, and Vinay Bhomia were study investigators. Each author contributed important intellectual content during manuscript drafting or revision and accepts accountability for the overall work by ensuring that questions pertaining to the accuracy or integrity of any portion of the work are appropriately investigated and resolved. All authors approved the final version of the manuscript for submission. [ZRC communication number: 660]

Funding Source

This trial was sponsored and funded by Cadila Healthcare Ltd., Ahmedabad, India.

Ethical approval

Written informed consent was obtained from all participants at the time of screening. This trial was initiated after obtaining the approvals of ECs, DCGI and registering the trial with the CTRI. This trial was conducted in accordance with the applicable local regulations.

Declaration of Competing Interest

All authors declared no competing interests.

References

- Bastard P, Rosen LB, Zhang Q, et al. Autoantibodies against type I IFNs in patients with life-threatening COVID-19. *Science*. 2020;370(6515):eabd4585. doi: 10.1126/science.abd4585. Epub 2020 Sep 24.
- Blanco-Melo D, Nilsson-Payant BE, Liu WC, et al. Imbalanced host response to SARS-CoV-2 drives development of COVID-19. *Cell*. 2020;181(5):1036-1045.e9. doi: 10.1016/j.cell.2020.04.026.
- Cevik M, Tate M, Lloyd O, et al. SARS-CoV-2, SARS-CoV, and MERS-CoV viral load dynamics, duration of viral shedding, and infectiousness: a systematic review and meta-analysis. *The Lancet Microbe*. 2020;2(1):e13-e22. doi :[https://doi.org/10.1016/S2666-5247\(20\)30172-5](https://doi.org/10.1016/S2666-5247(20)30172-5)
- Chakraborty C, Sharma AR, Bhattacharya M, et al. The 2019 novel coronavirus disease (COVID-19) pandemic: a zoonotic prospective. *Asian Pacific Journal of Tropical Medicine*. 2020;13(6):242. doi: 10.4103/1995-7645.281613
- Cucinotta D, Vanelli M. WHO declares COVID-19 a pandemic. *Acta Biomed.* 2020;91(1):157-60. doi: 10.23750/abm.v91i1.9397
- Falzarano D, De Wit E, Rasmussen AL, et al. Treatment with interferon- α 2b and ribavirin improves outcome in MERS-CoV–infected rhesus macaques. *Nat Med*. 2013;19(10):1313-17. doi: 10.1038/nm.3362. Epub 2013 Sep 8.
- Hadjadj J, Yatim N, Barnabei L, et al. Impaired type I interferon activity and inflammatory responses in severe Covid-19 patients. *Science*. 2020;369(6504):718-724. doi: 10.1126/science.abc6027. Epub 2020 Jul 13.

Lindsay KL, Trepo C, Heintges T, et al. A randomized, double-blind trial comparing pegylated interferon alfa-2b to interferon alfa-2b as initial treatment for chronic hepatitis C. *Hepatology*. 2001;34(2):395-403. doi: 10.1053/jhep.2001.26371

Lokugamage K, Hage A, Vries M, et al. Type I interferon susceptibility distinguishes SARS-CoV-2 from SARS-CoV. *J Virol*. 2020;94(23):e01410-20. doi: 10.1128/JVI.01410-20.

Loutfy MR, Blatt LM, Siminovitch KA, et al. Interferon alfacon-1 plus corticosteroids in severe acute respiratory syndrome: a preliminary study. *Jama*. 2003;290(24):3222-8. doi: 10.1001/jama.290.24.3222.

Ministry of Health and family welfare. Government of India. Clinical Management Protocol: COVID-19. 03 July 2020.

<https://www.mohfw.gov.in/pdf/UpdatedClinicalManagementProtocolforCOVID19dated03072020.pdf>

Nezhad FS, Mosaddeghi P, Negahdaripour M, et.al, Therapeutic approaches for COVID-19 based on the dynamics of interferon mediated immune responses, Preprints 2020, 2020030206 (doi: 10.20944/preprints202003.0206.v1).

Perlman S, Dandekar AA. Immunopathogenesis of Coronavirus Infections: Implications for SARS. *Nat Rev Immunol*. 2005;5(12):917-27. doi: 10.1038/nri1732.

Poynard T, McHutchison J, Manns M, et al. Impact of pegylated interferon alfa-2b and ribavirin on liver fibrosis in patients with chronic hepatitis C. *Gastroenterology*. 2002;122(5):1303-13. doi: <https://doi.org/10.1053/gast.2002.33023>

Schiffert JT, Johnston C, Wald A, et al. An early test-and-treat strategy for severe acute respiratory syndrome coronavirus 2. *Open Forum Infect Dis.* 2020;7(7):ofaa232. doi: 10.1093/ofid/ofaa232.

Spinelli A, Pellino G. COVID-19 pandemic: perspectives on an unfolding crisis. *Br J Surg.* 2020;107(7):785-787. doi: 10.1002/bjs.11627. Epub 2020 Mar 23.

Spinner CD, Gottlieb RL, Criner GJ, et al. Effect of remdesivir vs standard care on clinical status at 11 days in patients with moderate COVID-19: a randomized clinical trial. *Jama.* 2020;324(11):1048-57. doi: 10.1001/jama.2020.16349.

Stroher U, DiCaro A, Li Y, et al. Severe acute respiratory syndrome-related coronavirus is inhibited by interferon- α . *J Infect Dis.* 2004;189(7):1164-7. doi: 10.1086/382597. Epub 2004 Mar 12.

Vabret N, Britton GJ, Gruber C, et al. Immunology of COVID-19: current state of the science. *Immunity.* 2020;52(6): 910-41. doi: 10.1016/j.immuni.2020.05.002

Wang BX, Fish EN. Global virus outbreaks: interferons as 1st responders. *Semin Immunol.* 2019;43:101300. doi: 10.1016/j.smim.2019.101300

World Health Organization. COVID-19 weekly epidemiological update. 02 February 2021.

Zhang JJ, Dong X, Cao YY, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy.* 2020;75(7):1730-1741. doi: 10.1111/all.14238. Epub 2020 Feb 27.

Zhang Q, Bastard P, Liu Z, et al. Inborn errors of type I IFN immunity in patients with life-threatening COVID-19. *Science*. 2020;370(6515):eabd4570. doi: 10.1126/science.abd4570.

Zhao J, Yuan Q, Wang H, et al. Antibody responses to SARS-CoV-2 in patients of novel coronavirus disease 2019. *Clin Infect Dis*. 2020;71(16):2027-34. doi: 10.1093/cid/ciaa344.

Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395(10229):1054-62. doi: 10.1016/S0140-6736(20)30566-3. Epub 2020 Mar 11.

Zhou Q, Chen V, Shannon CP, et al. Interferon- α 2b Treatment for COVID-19. *Front Immunol*. 2020;11:1061. doi: 10.3389/fimmu.2020.01061

Table 1: Summary of Demographic Characteristics (Safety Population)

Subject Characteristics	PEG IFN- α2b + SOC (N=20)	SOC (N=20)	Total (N=40)
Age (Years), Mean (SD)	49.35 (14.89)	49.10 (12.44)	49.23 (13.55)
Sex, n (%)			
Female	9 (45.0%)	1 (5.0%)	10 (25.0%)
Male	11 (55.0%)	19 (95.0%)	30 (75.0%)
Height (cm), Mean (SD)	167.14 (9.34)	165.94 (8.46)	166.54 (8.82)
Weight (Kg), Mean (SD)	72.65 (12.94)	71.90 (11.34)	72.28 (12.01)
BMI (kg/m ²), Mean (SD)	26.12 (5.01)	26.21 (4.51)	26.16 (4.70)
<p>BMI = body mass index; N = number of subjects in treatment group; n = number of subjects in specified category; PEG IFN-α2b = pegylated interferon alfa-2b; SD = standard deviation; SOC = standard of Care</p>			

Table 2: Analysis of Proportion of Subjects With Clinical Improvement (Clinical Status) From Day 0 To Day 15, Measured Using the WHO 7-Point Ordinal Scale

Visit	Improvement	PEG IFN- α 2b + SOC (N=20)	SOC (N=19 ^a)	p-value ^b
Checkout Day 15	Yes	19 (95.00%)	13 (68.42%)	0.0436
	No	1 (5.00%)	6 (31.58%)	

N = number of subjects in treatment group; PEG IFN- α 2b = pegylated interferon alfa-2b; SOC = standard of Care

^a One subject was excluded from analysis because no post-baseline clinical status data available.

^b Fisher exact test has been used to calculate p-value.

P-value < 0.05 was considered as statistically significant

Table 3: Analysis of Change in Score From Day 0 To Day 15, Measured Using the WHO 7-Point Ordinal Scale

Change in score	PEG IFN- α 2b + SOC (N=20)	SOC (N=19 ^a)
Mean (SD)	-2.25 (0.55)	-2.05 (0.85)
Median	-2.00	-2.00
Min, Max	(-3, -1)	(-3, -1)
p-value ^b	<.0001	<.0001

N = number of subjects in treatment group; PEG IFN- α 2b = pegylated interferon alfa-2b;

SOC = standard of care

^a One subject was excluded from analysis because no post-baseline clinical status data available.


^b P-value has been calculated using Wilcoxon Signed Rank Test.

P-value <0.05 was considered as statistically significant.

Journal Pre-proof

Table 4: Summary of Adverse Events (Safety Population)


Preferred Term	PEG IFN-α2b + SOC (N=20)	SOC (N=20)
Adverse Event	11 (55.0%)	8 (40.0%)
Breathlessness	1 (5.0%)	3 (15.0%)
Chest Pain	0 (0%)	2 (10.0%)
Difficulty In Breathing	2 (10.0%)	2 (10.0%)
Dryness In Mouth	1 (5.0%)	1 (5.0%)
Headache	8 (40.0%)	4 (20.0%)
Hypoxia	1 (5.0%)	0 (0%)
Mouth Dry	0 (0%)	2 (10.0%)
Nausea	1 (5.0%)	1 (5.0%)
Vomiting	6 (30.0%)	2 (10.0%)
N = number of subjects in treatment group; PEG IFN- α 2b = pegylated interferon alfa-2b; SOC = standard of care		

Figure 1: Subject disposition

mITT = modified intent-to-treat, PEG IFN- α 2b = pegylated interferon alfa-2b, SOC = standard of care,


Figure 2: Qualitative RT-PCR for SARS-CoV-2 in pharyngeal swab

Number of SARS-CoV-2 positive subjects before and after treatment with PEG IFN- α 2b + SOC and SOC



PEG IFN- α 2b = pegylated interferon alfa-2b, SOC = standard of care

Figure 3: Occurrence and duration of supplemental oxygen



PEG IFN-α2b = pegylated interferon alfa-2b, SOC = standard of care