

COVID-19 and immunomodulator/ immunosuppressant use in dermatology



To the Editor: We read with great interest the Commentary by Lebwohl et al¹ recently published in the *Journal of the American Academy of Dermatology*. The authors provided a pertinent overview of infection risk associated with commonly used biologics to treat psoriasis in light of the current coronavirus disease 2019 (COVID-19) outbreak. We agree that this time has been particularly concerning for patients taking immunomodulators/immunosuppressants who are unsure of their risk for severe disease.

In response to the previous commentary, the goal of this letter is to expand and provide the latest information about COVID-19 along with considerations for addressing patient concerns surrounding dermatology-related immunomodulator/immunosuppressant use.

Theoretical data from previous coronavirus outbreaks has suggested a strong role for type I interferon, B-cell–released antibodies, tumor necrosis factor- α , and other cytokines in the viral immune response (Fig 1).^{2–4} Interleukin (IL) 17 cytokines are important for immune cell recruitment to infection sites to promote clearance, while also activating downstream cascades of cytokines and chemokines.⁴ IL-1 promotes fever and the differentiation of T-helper cells to IL-17–producing T cells. Tumor necrosis factor- α promotes dendritic cell differentiation, leukocyte recruitment, and mediates fever.⁴ Antibodies produced by plasma cells help to neutralize the virus, limit infection, and prevent future infections. Disruption of B-cell differentiation into plasma cells could limit antibody production.³

Broad immunosuppression across multiple cytokine axes with immunosuppressants has the potential to increase susceptibility, persistence, and reactivation of viral infections. Immunosuppressants decrease cytokines that recruit and differentiate immune cells needed to clear the infection. In addition, inflammatory mediators can become hyperactivated, resulting in a “cytokine storm,” which is the primary cause of death in severe disease.³

Whether withdrawal of broadly immunosuppressive therapies may increase the risk of precipitating cytokine storm is unknown. Therefore, classic immunosuppressants may present the most concerning risk for those affected by COVID-19 (Table D). Immunomodulators, such as biologics, that do not target vital domains within the viral immune

response may dampen, but not significantly affect viral clearance.

Currently, there are no data describing the benefits or risks of stopping immunomodulators/immunosuppressants during the COVID-19 outbreak. However, each medication’s mechanism of action, administration method/frequency, and pharmacokinetics/pharmacodynamics are important to consider. Nonbiologic medications, including small molecule inhibitors and immunosuppressants, are typically easier to stop and restart within days to weeks due to shorter half-life. Meanwhile, biologics generally have a longer half-life and include a risk of antidrug antibody formation with treatment cessation and subsequent continuation. However, biologics also tend to be more targeted and less involved in the previously mentioned components of the viral immune response. General medication considerations are included in Table I. Although patient dependent, clinicians may consider weaning patients with stable disease off of immunosuppressants.

Shared decision making is needed when deciding on a treatment plan that includes immunomodulators/immunosuppressants during the COVID-19 outbreak. Patients with existing comorbidities may require more conservative measures.⁵ Physicians should continue to consult with the Centers for Disease Control and Prevention Information for Healthcare Providers, which is updated daily (<https://www.cdc.gov/coronavirus/2019-nCoV/hcp/index.html>). Once again, we thank the authors for raising awareness of patient concerns during this evolving outbreak.

Kyla N. Price, BS,^a John W. Frew, MD,^b Jennifer L. Hsiao, MD,^c and Vivian Y. Shi, MD^d

From the College of Medicine, University of Illinois at Chicago, Chicago, Illinois^a; the Laboratory of Investigative Dermatology, The Rockefeller University, New York, New York^b; the Division of Dermatology, Department of Medicine, University of California, Los Angeles, Los Angeles, California^c; and the Division of Dermatology, Department of Medicine, University of Arizona, Tucson, Arizona.^d

Funding sources: None.

Conflicts of interest: Dr Shi is a stock shareholder of Learn Health and has served as an advisory board member, investigator, and/or received research funding from Sanofi Genzyme, Regeneron, AbbVie, Eli Lilly, Novartis, Sun Pharma, LEO Pharma, Pfizer, Menlo Therapeutics, Burt’s




Fig 1. COVID-19 viral immune response and targets of common dermatologic immunomodulators and immunosuppressants. **(Left)** (1) Person-to-person transmission of COVID-19 occurs through direct contact with respiratory secretions of infected individuals.² The virus invades host cells by binding to their receptors and fusing with the cell membrane. (2) It is hypothesized that once inside the body, the lung epithelial cells become the primary target, where the receptor binding domain of the virus spikes bind to angiotensin-converting enzyme 2 (ACE2) receptors of ACE2-expressing target cells. (3) Although not confirmed, it is believed the virus dampens the initial type 1 interferon (IFN) responses, which contributes to uncontrolled viral replication. (4) Once the virus is identified, macrophages present viral components to activate and induce (5) differentiation of T cells and B cells. (6) Activated B cells differentiate into plasma cells that produce antibodies important for neutralizing viruses. (7) The resulting inflammatory cytokines and antibodies continue to stimulate the production of additional cytokines and antibodies, which may contribute to the "cytokine storm" noted in those with severe disease. (8) The inflammatory cytokines and antibodies also promote the influx of neutrophils, monocytes, and macrophages along with additional inflammatory cytokines. **(Right)** The drug targets for common dermatologic immunomodulators and immunosuppressants have also been included in this diagram. *FGF*, Basic fibroblast growth factor; *G-CSF*, granulocyte-colony stimulating factor; *GM-CSF*, granulocyte-macrophage colony-stimulating factor; *IL*, interleukin; *IP10*, interferon γ -induced protein 10; *IRF*, interferon regulatory protein 1- α ; *NFAT*, nuclear factor of activated T cells; *NF- κ B*, nuclear factor- κ B; *PDE4*, phosphodiesterase 4; *PDGF*, platelet-derived growth factor; *PKA*, protein kinase A; *T_H*, T-helper cell; *TNF*, tumor necrosis factor; *VEGFA*, vascular endothelial growth factor A. Created with Biorender.com.

Table I. Considerations for commonly used immunomodulators and immunosuppressants for dermatologic conditions

Drug class	Mechanism of action	Drug name	Risk	Comments/considerations*		
Classic immunosuppressants						
Antimetabolites	Inhibits NF- κ B Calcineurin inhibitor	Corticosteroids	Likely concerning risk	Consider stopping when viral symptoms present especially with known or potential exposure		
		Tacrolimus				
		Cyclosporine				
		Mycophenolate mofetil				
		Azathioprine				
		Methotrexate				
Immunomodulators						
Monoclonal antibodies	TNF- α inhibition	Infliximab	Likely moderate risk	Continue if viral symptoms are mild, consider stopping if viral symptoms worsen or high fever develops		
Receptor fusion protein		Etanercept				
Monoclonal antibodies		Certolizumab				
Monoclonal antibodies		Adalimumab				
IL receptor modulators		Anakinra (IL-1)				
Monoclonal antibodies	IL inhibition	Dupilumab (IL-4)	Likely low risk	Continue unless severe symptoms present		
Monoclonal antibodies		Brodalumab (IL-17)			Likely moderate risk	Continue if viral symptoms are mild, consider stopping if viral symptoms worsen or high fever develops
Monoclonal antibodies		Secukinumab (IL-17a)				
Monoclonal antibodies		Ixekizumab (IL-17a)				
Monoclonal antibodies		Ustekinumab (IL-12/23)				
Monoclonal antibodies	Guselkumab (IL-23)					
Monoclonal antibodies	Anti-CD20 antibody	Rituximab	Likely concerning risk	Consider stopping when viral symptoms present especially with known or potential exposure.		
	PDE4 inhibition	Apremilast	Likely low risk	Continue unless severe symptoms present		

IL, Interleukin; NF- κ B, nuclear factor κ B; PDE4, phosphodiesterase 4.

*General considerations only, medication use should be considered based on each individual patient's risk and disease profile.

Bees, GpSkin, Altus Labs, and Skin Actives Scientific. There were no incentives or transactions, financial or otherwise, relevant to this manuscript. Kyla N. Price and Drs Frew and Hsiao have no conflicts of interest to declare.

IRB approval status: Not applicable.

Accepted for publication March 20, 2020.

Reprints not available from the authors.

Correspondence to: Vivian Y. Shi, MD, Dermatology Division, University of Arizona, 7165 N Pima Canyon Dr, Tucson, AZ 85718

E-mail: vivian.shi.publications@gmail.com

REFERENCES

1. Lebwohl M, Rivera-Oyola R, Murrell DF. Should biologics for psoriasis be interrupted in the era of COVID-19? *J Am Acad Dermatol*. 2020;82(5):1217-1218.
2. Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J Autoimmun*. 2020;102433.
3. Prompetchara E, Ketloy C, Palaga T. Immune responses in COVID-19 and potential vaccines: lessons learned from SARS and MERS epidemic. *Asian Pac J Allergy Immunol*. 2020;38(1):1-9.
4. Li G, Fan Y, Lai Y, et al. Coronavirus infections and immune responses. *J Med Virol*. 2020;92(4):424-432.
5. Centers for Disease Control and Prevention. Coronavirus (COVID-19). Available at: <https://www.cdc.gov/coronavirus/2019-ncov/index.html>. Accessed March 9, 2020.

<https://doi.org/10.1016/j.jaad.2020.03.046>