

Antibody therapies for the treatment of COVID-19

Zhiqiang Ku, Xiaohua Ye, Georgina To'a Salazar, Ningyan Zhang and Zhiqiang An*

Texas Therapeutics Institute, Brown Foundation Institute of Molecular Medicine,
McGovern Medical School, The University of Texas Health Science Center at Houston,
Houston, Texas 77030, USA

*Corresponding Author

Zhiqiang An

E-mail: Zhiqiang.An@uth.tmc.edu

Keywords: COVID-19, SARS-CoV-2, anti-viral therapy, convalescent plasma therapy, spike protein, neutralizing antibody

Significance

Development of SARS-CoV-2 neutralizing antibodies with the desired efficacy and safety profile is a critical part of the toolbox of therapies for the treatment of COVID-19. We discuss in this review the current state of discovery and development of such antibodies.

© The Author(s) 2020. Published by Oxford University Press on behalf of Antibody Therapeutics. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com

Abstract

An outbreak of COVID-19, the disease caused by infection of the coronavirus SARS-CoV-2, that began in December 2019 in Wuhan, China has caused more than 2,990,559 confirmed human infections and 207,446 deaths as of April 27, 2020 (Coronavirus COVID-19 Global Cases by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University). Scientists are working quickly on multiple aspects of the pandemic. Genetic analyses are conducted to reveal the source and evolution of SARS-CoV-2, providing knowledge that can be used to contain it and to avoid future outbreaks. Epidemiological studies which incorporates lessons learned from outbreaks of previous related viral diseases can guide development of public health measures effective to contain the current and future outbreaks. Basic virology studies reveal viral structure and function. Pathology studies inform development of strategies to interfere with infection. COVID-19 prevention and treatment strategies are being developed in preclinical and clinical studies. Antibody-based therapy is one viable treatment option. Here, we discuss some of the most active areas of developing strategies to treat COVID-19, focusing on approaches to generate neutralizing antibodies against SARS-CoV-2 for prophylactic and therapeutic treatment of COVID-19.

A. The structures and functions of SARS-CoV-2 viral proteins

SARS-CoV-2, along with SARS-CoV and MERS-CoV, belongs to the *Betacoronavirus* genus in the Coronaviridae family (Figure 1A)^{1,2}. Bats are possible origins of SARS-CoV-2. SARS-CoV-2 is likely a result of natural selection either in an animal host before zoonotic transfer or in humans following zoonotic transfer^{2,3}. Similar to its family members SARS-CoV and MERS-CoV, SARS-CoV-2 has a large, positive-sense RNA genome, which encodes 4 structural proteins, 16 nonstructural proteins (nsp1-16), and some accessory

proteins (Figure 1B)^{1,2}. The structural proteins are spike (S), envelope (E), membrane (M), and nucleocapsid (N) proteins⁴. The spike protein comprises an N-terminal S1 subunit responsible for receptor binding and a C-terminal S2 subunit responsible for membrane fusion (Figure 1C)⁵. The S1 subunit is further divided into the N-terminal domain (NTD), the receptor-binding domain (RBD), subdomain 1 (SD1) and subdomain 2 (SD2). The S2 subunit is further divided into the fusion peptide (FP), the heptad repeat 1 (HR1) and heptad repeat 2 (HR2) (Figure 1C)⁶. Like SARS-CoV, SARS-CoV-2 enters cells through binding of the host cellular receptor angiotensin-converting enzyme 2 (ACE2) via its spike protein². In contrast, MERS-CoV enters cells through binding of the host receptor dipeptidyl peptidase 4 (DPP4) via its spike protein⁷. These host receptors are indispensable for virus infection^{2,4}. Receptor binding triggers a conformational change of the spike protein to an activated state⁸. The activated spike is cleaved by a protease (TMPRSS2 for SARS-CoV and SARS-CoV-2) at the S1/S2 site, releasing the S1 subunit and exposing the FP on the S2 subunit⁹. The FP inserts into target cell membrane, HR1 and HR2 refold to form a postfusion conformation that drives viral membrane fusion with target cells (Figure 1D)^{8,10}. The cryo-EM structure of the SARS-CoV-2 spike in the prefusion conformation was recently published¹¹. The interaction between SARS-CoV-2 spike RBD and the full-length human ACE2 has also been revealed by cryo-EM or crystallization^{6,12-14}. Similar to the spike protein of SARS-CoV, the SARS-CoV-2 spike protein also possesses extensive glycosylation, which may be important for virus binding to cells and facilitate immune evasion through epitope masking^{15,16}. The non-structural proteins encoded by coronaviruses are essential for virus replication inside cells¹. For example, the RNA-dependent RNA polymerase (RdRp, also known as nsp12) is the key component of coronaviral replication and transcription machinery¹. The cryo-EM structure

of RdRp in complex with cofactors nsp7 and nsp8 was recently published¹⁷. The rapid advance in the basic knowledge of SARS-CoV-2 paves the way for the discovery and development of vaccines and therapeutics for the prevention and treatment of COVID-19.

B. Repurposing drugs for the treatment of COVID-19

Discovery and development of drugs and vaccines against COVID-19 is in full swing in academic and biopharmaceutical company laboratories worldwide. Because of the urgency of finding treatments for the current pandemic, existing drugs developed for other viral diseases are being tested in the clinic for the treatment of COVID-19. Nucleoside analogues in the form of adenine or guanine derivatives target RdRp and block viral RNA synthesis in a broad spectrum of RNA viruses. Both approved nucleoside analogues, such as favipiravir and ribavirin, and experimental nucleoside analogues, such as remdesivir and galidesivir, may block SARS-CoV-2 RdRp and protect against COVID-19¹⁸ (Figure 1D). However, the coronavirus protein nsp14 has a unique exoribonuclease (ExoN) function, which provides the proofreading capability. This mechanism may affect the susceptibility of coronaviruses to nucleoside analogues¹⁹. Approved viral protease inhibitors, such as the HIV aspartic protease inhibitors lopinavir and ritonavir, have shown efficacy in a clinical trial²⁰. It is hypothesized that lopinavir and ritonavir exhibit their efficacy by inhibiting the 3-chymotrypsin-like protease²⁰ (Figure 1D).

In addition to repurposing existing antiviral drugs, drugs used to modulate the host immune system for other disease indications are also being tested in the clinic for the treatment of COVID-19. The antimalarial and anti-inflammatory drugs hydroxychloroquine and chloroquine showed *in vitro* activity against SARS-CoV-2²¹. Hydroxychloroquine and chloroquine may exhibit efficacy against COVID-19 by modulation of immune system and by affecting cell entry and replication of SARS-CoV-2 through pH modulation and

interfering post-translational modifications of viral proteins or receptors²². Adjunctive and supportive therapies such as azithromycin and tocilizumab are also being clinically tested for the treatment of COVID-19. Azithromycin is a macrolide antibacterial that may prevent bacterial superinfection on COVID-19 patients; macrolides may have immunomodulatory properties in pulmonary inflammatory disorders²³. Tocilizumab is an interleukin-6 (IL-6) receptor-blocking antibody that inhibits IL-6-mediated signaling for the treatment of inflammatory diseases^{24, 25}. The rationale for testing the efficacy of tocilizumab against COVID-19 is that IL-6 is a proinflammatory cytokine; cytokine release syndrome may be a component of severe disease in COVID-19 patients²⁶. Discussed here are a few examples of a vast array of drugs and drug combinations being tested for the treatment of COVID-19. Although some clinical trials have produced promising results, no drugs have been definitively validated for COVID-19 treatment to date. Repurposing existing drugs and *de novo* discovery of new drugs for the treatment of COVID-19 is an active on-going effort worldwide.

C. Vaccines for COVID-19

An effective vaccine to prevent SARS-CoV-2 infection is urgently needed to control the global pandemic of COVID-19. Unfortunately, there are no currently approved human coronavirus vaccines. The spike protein is a vaccine target for coronavirus; studies on SARS-CoV and MERS-CoV demonstrated that antibodies targeting the spike protein, especially its receptor binding domain (RBD), efficiently neutralize virus infection^{5, 27}.

Although SARS-CoV-2 and SARS-CoV share high sequence identity and use the same receptor for cell entry, the sera of patients who recovered from SARS-CoV and SARS-CoV-2 showed limited cross neutralization^{2, 28}. In a recent study, the SARS-CoV S murine polyclonal antibodies potently inhibited SARS CoV-2 S mediated entry into cells, indicating

that cross-neutralizing antibodies targeting conserved S epitopes can be elicited upon vaccination²⁹. Multiple biopharmaceutical companies and academic institutions worldwide have announced SARS-CoV-2 vaccine programs. These programs target major strategies for vaccine development, which include production of live attenuated whole virion vaccines, inactivated whole virion vaccines, spike protein targeted nucleic acid vaccines, recombinant protein vaccines, and viral vector-based vaccines³⁰⁻³². Currently, the most advanced SARS-CoV-2 vaccine candidates are in Phase I clinical trials.

D. Convalescent plasma therapy

Convalescent plasma (CP) from patients who recovered from a viral disease is a therapeutic tool used when no approved specific antiviral agents are available. During outbreaks of SARS-CoV, MERS-CoV, influenza A (including H7N9, H1N1, and H5N1), and Ebola, CP treatment of infected patients showed beneficial effects including relief of clinical symptoms and reduced mortality³³⁻³⁸. CP transfusion is being tested to rescue patients with severe COVID-19. In one trial, single doses of 200 ml CP with a >1:640 neutralizing antibody titer were administered to 10 patients with severe COVID-19 at a median time of 16.5 days from onset of illness³⁹. In patients who received CP transfusion, clinical symptoms (including fever, cough, shortness of breath, and chest pain) disappeared or largely improved. Viral RNA disappeared and there was a reduction of pulmonary lesions and reduced patient death compared to a matched historic control group³⁹. Importantly, no adverse events were reported among patients receiving CP transfusion. The safety profile and potential therapeutic effect of CP transfusion among patients with severe COVID-19 are also supported by other reports with fewer patients⁴⁰.⁴¹. Notably, in addition to CP transfusion, all patients received standard care, including antiviral therapy. This limits the use of these preliminary studies for the assessment of CP

transfusion as a COVID-19 therapy. Before this COVID-19 therapy can be widely used alone or in combination with other treatment(s), it must be fully assessed in a randomized controlled clinical trial in a larger patient cohort. A minimum effective neutralizing antibody titer and optimal treatment time point needs to be established for CP therapy. Although CP is readily accessible during the pandemic, major challenges remain for large-scale application of CP treatment. These challenges include the limited availability of CP in large quantities and donor dependent variability in antibody specificities and titers. The US FDA has approved the use of plasma from recovered patients to treat patients who are critically ill with COVID-19, provided that doctors get approval ⁴². The ongoing COVID-19 pandemic provides an unprecedented opportunity to perform clinical studies of CP treatment against a viral agent. If large-scale randomized clinical trials can demonstrate the efficacy and safety of CP, its therapeutic use among patients with severe disease and prophylactic use in high-risk populations could change the course of this ongoing pandemic.

E. SARS-CoV-2 neutralizing antibodies for the treatment of COVID-19

Neutralizing antibodies are an important component in host immune responses to viral pathogens. Neutralizing monoclonal antibodies have been developed as therapies for the treatment of viral infections including RSV, Influenza, Ebola, HIV, HCMV and Rabies ⁴³. The process for the discovery and development of viral neutralizing antibody therapies are well-established ⁴³. Scientists around the world have initiated programs to develop SARS-CoV-2 neutralizing antibodies. Currently, all published SARS-CoV-2 neutralizing antibodies are at early preclinical stages. We discuss here the current state of design and development of such antibodies.

1. *Viral targets for development of neutralizing antibodies against SARS-CoV-2 infection*

The spike protein of SARS-CoV-2 plays an essential role in virus entry into host cells and is a primary target of neutralizing antibodies^{5,9} (Figures 1C,D). Due to the functionality and high immunogenicity of the coronavirus S1 subunit, most neutralizing antibodies characterized for coronaviruses to date target S1, in particular the S1-RBD⁴⁴⁻⁴⁶. Two MERS-CoV neutralizing mAbs, G2 and 7D10, target the S1-NTD region and function by blocking spike protein interaction with the host receptor DPP4^{47,48}. Compared to the S1 subunit, the coronavirus S2 subunit is more conserved and bears epitopes that could potentially be targeted by broadly neutralizing antibodies^{10,49}. Generation of antibodies with broad neutralizing activity against different coronaviruses, or at least SARS-related coronaviruses, would be of great value for confronting future waves of coronavirus-related disease. However, broadly neutralizing antibodies against different human coronaviruses are very rare, probably related to the sequence variance of spike protein and cryptic nature of the highly conserved epitopes^{28,50}. The S2 conformation is highly dynamic during membrane fusion, presenting a major challenge in preparing antigens for discovery of antibodies against this protein¹⁰. Antigen stabilizing strategies used in discovery of antibodies against HIV and RSV proteins may be explored in the design of stable coronavirus S2 proteins⁵¹⁻⁵³.

2. Methods for isolating SARS-CoV-2 neutralizing antibodies

To target the whole spike protein or domain proteins, several well-established methods have been used for isolating SARS-CoV-2 monoclonal antibodies (Table 1). Xiong *et al.* isolated several neutralizing antibodies from mice immunized with SARS-CoV-2 RBD using the hybridoma technology⁵⁴. Wu *et al.* identified high-affinity human single-domain antibodies targeting five types of neutralizing or non-neutralizing epitopes on SARS-CoV-2 RBD through panning a phage-displayed single-domain antibody library⁵⁵. Ju *et al.*

characterized 206 RBD-specific mAbs isolated from single B cells of eight SARS-CoV-2 infected patients⁵⁶. Once a panel of monoclonal antibodies is generated, biochemical assays, such as ELISA binding, cell-based binding, affinity measurement, and epitope mapping and binning, are commonly used to characterize the antibodies (Table 1). Lead antibody candidates are then evaluated for their neutralization activities in *in vitro* cell based assays and in *in vivo* animal models.

To reduce the time from that required for *de novo* discovery of SARS-CoV-2 neutralizing antibodies, researchers have assessed available SARS-CoV antibodies for cross-neutralization of SARS-CoV-2, given the ~76% sequence identities in their spike proteins^{56,57}. For example, a SARS-CoV mAb (47D11) generated from a hybridoma of humanized mice neutralized SARS-CoV with an IC₅₀ of 0.19 µg/ml and cross neutralized SARS-CoV-2 with an IC₅₀ of 0.57 µg/ml. Although the 47D11 mAb binds to the RBD of both viruses, it could not block RBD binding to ACE2, suggesting presence of a broadly neutralizing epitope that is not directly involved in receptor binding⁵⁷. Another SARS-CoV mAb, CR3022, binds to the RBD of both SARS-CoV-2 and SARS-CoV, but only neutralizes SARS-CoV⁵⁰. A structural study indicates that CR3022 recognizes a highly conserved cryptic epitope on RBD that is distinct from the receptor-binding site⁵⁰.

3. *In vitro* assays for screening and selection of SARS-CoV-2 neutralizing antibodies

Antibodies that bind to the spike protein with high affinity can be first tested for their ability to block spike interaction with ACE2 or directly tested in cell based viral neutralization assays *in vitro*. Two *in vitro* assay systems are commonly used to evaluate the neutralization activity of antibodies against coronavirus⁵⁶. One is based on live virus infection of the Vero E6 cell line; it is the gold standard of *in vitro* neutralization assay.

Briefly, after infection by SARS-CoV-2, the cells can display cytopathic effects (CPE), which can be visually observed or form viral plaques that can be detected by immunofluorescent staining⁵⁶. However, this assay system is restricted, as it needs to be performed in Biosafety Level-3 (BSL-3) facilities. The other *in vitro* assay system commonly used to assess antibodies against coronavirus is based on a pseudovirus. The pseudovirus bears the spike of SARS-CoV2 in the lentiviral/retroviral backbones and has reporter genes such as GFP or luciferase. Expression of the reporter genes is used to indicate infection^{54, 56}. This method is much less restricted by the need of BSL-3 facilities, and can be high throughput. However, the conformation and number of the spike proteins on the pseudovirus may be different from those on the authentic virus. Often, researchers use the pseudovirus assay for initial high throughput screenings of lead antibodies followed by the live virus assay to further validate the lead candidates.

4. ***In vivo models for testing SARS-CoV-2 neutralizing antibodies***

Monkey and mouse models have been reported for testing SARS-CoV-2-neutralizing antibodies⁵⁸⁻⁶⁰. In the monkey study, researchers found that rhesus macaques infected with SARS-CoV-2 through the intratracheal route had mild illness, and their lungs showed signs of pneumonia similar to those in humans with COVID-19⁵⁸. After 3 or 6 days of infection, the virus could be isolated from bronchus, lung tissues, and oropharyngeal swabs. However, no monkey developed severe symptoms during the study⁵⁸. In a study using a mouse model, transgenic mice with human ACE2 expression, but not wild-type mice, were found to be susceptible to SARS-CoV-2 infection⁵⁹. Mice inoculated with SARS-CoV-2 through the intranasal route had a 5~10% loss of body weight and histopathology in the lung, such as interstitial pneumonia, and infiltration of immune cells. No mice died during the study⁵⁹. To establish a model that can mimic more severe human

infections, different animal models and other experimental factors must be considered and tested.

5. ADE consideration and Fc engineering

Antibody-dependent enhancement (ADE) is the phenomenon of non-neutralizing or sub-neutralizing antibodies facilitating virus infection, leading to more severe disease. The most widely known example is dengue ADE, which has various mechanisms^{61, 62}. In one mechanism, certain immune cells, which don't express the receptor for virus entry but have the Fcγ receptors (FcγR) on surface, can be infected by dengue virus through the FcγR mediated virus entry pathway⁶². Although it is unknown whether ADE happens in SARS-CoV-2 infection and COVID-19 disease, several ADE mechanisms have been described in other human coronaviruses, including the closely related SARS-CoV⁶³⁻⁶⁶. In one mechanism, passive transfer of anti-spike IgGs, which were elicited by vaccine immunization, failed to prevent infection and enhanced lung pathology in a monkey model of SARS-CoV infection⁶⁴. The antibodies exaggerated disease by modulating the monocyte and macrophage function in an FcγR dependent manner⁶⁴. In another mechanism revealed by two studies, the neutralizing mAb S230 for SARS-CoV and Mersmab1 for MERS-CoV, triggered membrane fusion via receptor mimicry when they bound to the RBD of the spike^{66, 67}. The presence of Mersmab1 enhanced the entry of MERS-CoV pseudovirus into macrophages and into FcγRII-expressing non-susceptible HEK293T cells⁶⁷. Notably, the SARS-CoV-2 serum has a substantial cross-reactivity to the spike from SARS-CoV and MERS-CoV, while the cross-neutralization activity is very limited⁵⁶. One possible explanation is that most antibodies target domains that are outside of the RBD and have few neutralizing epitopes⁵⁶. These studies suggest that ADE might occur in SARS-CoV-2 infections. Therefore, clinical application of convalescent plasma

therapy, design and development of vaccines and therapeutic antibodies for COVID-19 should take the risk of ADE into consideration.

Perspective

The unprecedented impact of the current COVID-19 pandemic on the human race calls for unprecedented response from all aspects of society to combat the disease. To address the immediate need for therapies, repurposing existing drugs and strategies is a logical first step. This includes testing antivirals, drugs used to modulate the host immune system, and CP transfusion as therapies for COVID-19. It is likely that SARS-CoV-2 will stay with us as a seasonal infection⁶⁸. Development of effective vaccines is the best weapon to confront COVID-19 if SARS-CoV-2 circulates as a seasonal virus or another coronavirus emerges as a serious threat in the future.

One challenge for developing COVID-19 therapies is the complexity of pathology and immunology associated with SARS-CoV-2 infection and its comorbidities. The vast majority of deaths have occurred in patients with underlying conditions⁶⁹. Considerations for designing clinical trials for SARS-CoV-2 therapies include optimization of dose and dosing frequency, disease stage at which to administer therapy, and combination with other therapies. For SARS-CoV-2 infection, virus shedding happens before onset of clinical symptoms and the viral dynamics in infected patients are still yet to be fully determined⁷⁰; treatment should be administered early enough to avoid severe symptoms. This timing of therapy can be achieved with a rapid, point of care diagnostic test.

Well-established antibody technologies enable laboratories in academia and drug development companies to isolate SARS-CoV-2 neutralizing antibodies in months or even weeks. However, development of SARS-CoV-2 neutralizing antibody therapies with the desired efficacy for human use is a complex process with many challenges. To best

choose epitopes, antibody formats, and antibody combinations, we must determine how fast SARS-CoV-2 can generate escape mutants and which epitopes are more vulnerable to neutralizing antibodies. Another important consideration is the engineering of Fc region of the neutralizing antibodies to avoid the potential antibody-dependent enhancement (ADE). Despite the challenges, development of neutralizing antibodies is a critical part of the toolbox of therapies for the treatment of COVID-19.

Acknowledgement

This work was supported in part by a Welch Foundation grant AU-0042-20030616 and Cancer Prevention and Research Institute of Texas (CPRIT) Grants RP150551 and RP190561.

Conflict of interest statement

The authors declare no potential conflicts of interest.

References

- [1] de Wit E, van Doremalen N, Falzarano D, Munster VJ: SARS and MERS: recent insights into emerging coronaviruses. *Nature reviews Microbiology* 2016, 14:523-34.
- [2] Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL, Chen HD, Chen J, Luo Y, Guo H, Jiang RD, Liu MQ, Chen Y, Shen XR, Wang X, Zheng XS, Zhao K, Chen QJ, Deng F, Liu LL, Yan B, Zhan FX, Wang YY, Xiao GF, Shi ZL: A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020, 579:270-3.
- [3] Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF: The proximal origin of SARS-CoV-2. *Nature Medicine* 2020.
- [4] Cui J, Li F, Shi ZL: Origin and evolution of pathogenic coronaviruses. *Nature reviews Microbiology* 2019, 17:181-92.
- [5] Du L, He Y, Zhou Y, Liu S, Zheng BJ, Jiang S: The spike protein of SARS-CoV--a target for vaccine and therapeutic development. *Nature reviews Microbiology* 2009, 7:226-36.
- [6] Lan J, Ge J, Yu J, Shan S, Zhou H, Fan S, Zhang Q, Shi X, Wang Q, Zhang L, Wang X: Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature* 2020.
- [7] Raj VS, Mou H, Smits SL, Dekkers DH, Muller MA, Dijkman R, Muth D, Demmers JA, Zaki A, Fouchier RA, Thiel V, Drosten C, Rottier PJ, Osterhaus AD, Bosch BJ, Haagmans BL: Dipeptidyl peptidase 4 is a functional receptor for the emerging human coronavirus-EMC. *Nature* 2013, 495:251-4.

- [8] Li F: Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annual review of virology* 2016, 3:237-61.
- [9] Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu NH, Nitsche A, Muller MA, Drosten C, Pohlmann S: SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 2020.
- [10] Walls AC, Tortorici MA, Snijder J, Xiong X, Bosch BJ, Rey FA, Velesler D: Tectonic conformational changes of a coronavirus spike glycoprotein promote membrane fusion. *Proceedings of the National Academy of Sciences of the United States of America* 2017, 114:11157-62.
- [11] Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, Graham BS, McLellan JS: Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 2020, 367:1260-3.
- [12] Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q: Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science* 2020, 367:1444-8.
- [13] Shang J, Ye G, Shi K, Wan Y, Luo C, Aihara H, Geng Q, Auerbach A, Li F: Structural basis of receptor recognition by SARS-CoV-2. *Nature* 2020.
- [14] Wang Q, Zhang Y, Wu L, Niu S, Song C, Zhang Z, Lu G, Qiao C, Hu Y, Yuen KY, Wang Q, Zhou H, Yan J, Qi J: Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2. *Cell* 2020.
- [15] Walls AC, Tortorici MA, Frenz B, Snijder J, Li W, Rey FA, DiMaio F, Bosch BJ, Velesler D: Glycan shield and epitope masking of a coronavirus spike protein observed by cryo-electron microscopy. *Nature structural & molecular biology* 2016, 23:899-905.
- [16] Vankadari N, Wilce JA: Emerging WuHan (COVID-19) coronavirus: glycan shield and structure prediction of spike glycoprotein and its interaction with human CD26. *Emerging microbes & infections* 2020, 9:601-4.
- [17] Gao Y, Yan L, Huang Y, Liu F, Zhao Y, Cao L, Wang T, Sun Q, Ming Z, Zhang L, Ge J, Zheng L, Zhang Y, Wang H, Zhu Y, Zhu C, Hu T, Hua T, Zhang B, Yang X, Li J, Yang H, Liu Z, Xu W, Guddat LW, Wang Q, Lou Z, Rao Z: Structure of the RNA-dependent RNA polymerase from COVID-19 virus. *Science* 2020.
- [18] Costanzo M, De Giglio MAR, Roviello GN: SARS CoV-2: Recent Reports on Antiviral Therapies Based on Lopinavir/Ritonavir, Darunavir/Umifenovir, Hydroxychloroquine, Remdesivir, Favipiravir and Other Drugs for the Treatment of the New Coronavirus. *Current medicinal chemistry* 2020.
- [19] Agostini ML, Andres EL, Sims AC, Graham RL, Sheahan TP, Lu X, Smith EC, Case JB, Feng JY, Jordan R, Ray AS, Cihlar T, Siegel D, Mackman RL, Clarke MO, Baric RS, Denison MR: Coronavirus Susceptibility to the Antiviral Remdesivir (GS-5734) Is Mediated by the Viral Polymerase and the Proofreading Exoribonuclease. *mBio* 2018, 9.
- [20] Zumla A, Chan JF, Azhar EI, Hui DS, Yuen KY: Coronaviruses - drug discovery and therapeutic options. *Nature reviews Drug discovery* 2016, 15:327-47.
- [21] Yao X, Ye F, Zhang M, Cui C, Huang B, Niu P, Liu X, Zhao L, Dong E, Song C, Zhan S, Lu R, Li H, Tan W, Liu D: In Vitro Antiviral Activity and Projection of Optimized Dosing Design of Hydroxychloroquine for the Treatment of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2020.
- [22] Devaux CA, Rolain JM, Colson P, Raoult D: New insights on the antiviral effects of chloroquine against coronavirus: what to expect for COVID-19? *International journal of antimicrobial agents* 2020:105938.

- [23] Gautret P, Lagier JC, Parola P, Hoang VT, Meddeb L, Mailhe M, Doudier B, Courjon J, Giordanengo V, Vieira VE, Dupont HT, Honore S, Colson P, Chabriere E, La Scola B, Rolain JM, Brouqui P, Raoult D: Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *International journal of antimicrobial agents* 2020:105949.
- [24] Sheppard M, Laskou F, Stapleton PP, Hadavi S, Dasgupta B: Tocilizumab (Actemra). *Human vaccines & immunotherapeutics* 2017, 13:1972-88.
- [25] Paul-Pletzer K: Tocilizumab: blockade of interleukin-6 signaling pathway as a therapeutic strategy for inflammatory disorders. *Drugs Today (Barc)* 2006, 42:559-76.
- [26] Slater H: FDA Approves Phase III Clinical Trial of Tocilizumab for COVID-19 Pneumonia. <https://www.cancernetwork.com/news/fda-approves-phase-iii-clinical-trial-tocilizumab-covid-19-pneumonia>, 2020.
- [27] Wang Q, Wong G, Lu G, Yan J, Gao GF: MERS-CoV spike protein: Targets for vaccines and therapeutics. *Antiviral research* 2016, 133:165-77.
- [28] Ou X, Liu Y, Lei X, Li P, Mi D, Ren L, Guo L, Guo R, Chen T, Hu J, Xiang Z, Mu Z, Chen X, Chen J, Hu K, Jin Q, Wang J, Qian Z: Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. *Nature communications* 2020, 11:1620.
- [29] Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D: Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* 2020, 181:281-92 e6.
- [30] Amanat F, Krammer F: SARS-CoV-2 Vaccines: Status Report. *Immunity* 2020.
- [31] Chen WH, Strych U, Hotez PJ, Bottazzi ME: The SARS-CoV-2 Vaccine Pipeline: an Overview. *Current tropical medicine reports* 2020:1-4.
- [32] Thanh Le T, Andreadakis Z, Kumar A, Gomez Roman R, Tollefsen S, Saville M, Mayhew S: The COVID-19 vaccine development landscape. *Nature reviews Drug discovery* 2020.
- [33] van Griensven J, Edwards T, de Lamballerie X, Semple MG, Gallian P, Baize S, Horby PW, Raoul H, Magassouba N, Antierens A, Lomas C, Faye O, Sall AA, Fransen K, Buyze J, Ravinetto R, Tiberghien P, Claeys Y, De Crop M, Lynen L, Bah EI, Smith PG, Delamou A, De Weggheleire A, Haba N, Ebola-Tx C: Evaluation of Convalescent Plasma for Ebola Virus Disease in Guinea. *The New England journal of medicine* 2016, 374:33-42.
- [34] Cheng Y, Wong R, Soo YO, Wong WS, Lee CK, Ng MH, Chan P, Wong KC, Leung CB, Cheng G: Use of convalescent plasma therapy in SARS patients in Hong Kong. *European journal of clinical microbiology & infectious diseases* : official publication of the European Society of Clinical Microbiology 2005, 24:44-6.
- [35] Wu XX, Gao HN, Wu HB, Peng XM, Ou HL, Li LJ: Successful treatment of avian-origin influenza A (H7N9) infection using convalescent plasma. *International journal of infectious diseases* : IJID : official publication of the International Society for Infectious Diseases 2015, 41:3-5.
- [36] Hung IF, To KK, Lee CK, Lee KL, Chan K, Yan WW, Liu R, Watt CL, Chan WM, Lai KY, Koo CK, Buckley T, Chow FL, Wong KK, Chan HS, Ching CK, Tang BS, Lau CC, Li IW, Liu SH, Chan KH, Lin CK, Yuen KY: Convalescent plasma treatment reduced mortality in patients with severe pandemic influenza A (H1N1) 2009 virus infection. *Clinical infectious diseases* : an official publication of the Infectious Diseases Society of America 2011, 52:447-56.
- [37] Zhou B, Zhong N, Guan Y: Treatment with convalescent plasma for influenza A (H5N1) infection. *The New England journal of medicine* 2007, 357:1450-1.


- [38] Ko JH, Seok H, Cho SY, Ha YE, Baek JY, Kim SH, Kim YJ, Park JK, Chung CR, Kang ES, Cho D, Muller MA, Drosten C, Kang CI, Chung DR, Song JH, Peck KR: Challenges of convalescent plasma infusion therapy in Middle East respiratory coronavirus infection: a single centre experience. *Antiviral therapy* 2018, 23:617-22.
- [39] Duan K, Liu B, Li C, Zhang H, Yu T, Qu J, Zhou M, Chen L, Meng S, Hu Y, Peng C, Yuan M, Huang J, Wang Z, Yu J, Gao X, Wang D, Yu X, Li L, Zhang J, Wu X, Li B, Xu Y, Chen W, Peng Y, Hu Y, Lin L, Liu X, Huang S, Zhou Z, Zhang L, Wang Y, Zhang Z, Deng K, Xia Z, Gong Q, Zhang W, Zheng X, Liu Y, Yang H, Zhou D, Yu D, Hou J, Shi Z, Chen S, Chen Z, Zhang X, Yang X: Effectiveness of convalescent plasma therapy in severe COVID-19 patients. *Proceedings of the National Academy of Sciences of the United States of America* 2020.
- [40] Shen C, Wang Z, Zhao F, Yang Y, Li J, Yuan J, Wang F, Li D, Yang M, Xing L, Wei J, Xiao H, Yang Y, Qu J, Qing L, Chen L, Xu Z, Peng L, Li Y, Zheng H, Chen F, Huang K, Jiang Y, Liu D, Zhang Z, Liu Y, Liu L: Treatment of 5 Critically Ill Patients With COVID-19 With Convalescent Plasma. *Jama* 2020.
- [41] Zhang B, Liu S, Tan T, Huang W, Dong Y, Chen L, Chen Q, Zhang L, Zhong Q, Zhang X, Zou Y, Zhang S: Treatment with convalescent plasma for critically ill patients with SARS-CoV-2 infection. *Chest* 2020.
- [42] Tanne JH: Covid-19: FDA approves use of convalescent plasma to treat critically ill patients. *Bmj* 2020, 368:m1256.
- [43] Salazar G, Zhang N, Fu TM, An Z: Antibody therapies for the prevention and treatment of viral infections. *NPJ vaccines* 2017, 2:19.
- [44] Jiang S, Hillyer C, Du L: Neutralizing Antibodies against SARS-CoV-2 and Other Human Coronaviruses. *Trends in immunology* 2020.
- [45] Zhu Z, Chakraborti S, He Y, Roberts A, Sheahan T, Xiao X, Hensley LE, Prabakaran P, Rockx B, Sidorov IA, Corti D, Vogel L, Feng Y, Kim JO, Wang LF, Baric R, Lanzavecchia A, Curtis KM, Nabel GJ, Subbarao K, Jiang S, Dimitrov DS: Potent cross-reactive neutralization of SARS coronavirus isolates by human monoclonal antibodies. *Proceedings of the National Academy of Sciences of the United States of America* 2007, 104:12123-8.
- [46] Ying T, Du L, Ju TW, Prabakaran P, Lau CC, Lu L, Liu Q, Wang L, Feng Y, Wang Y, Zheng BJ, Yuen KY, Jiang S, Dimitrov DS: Exceptionally potent neutralization of Middle East respiratory syndrome coronavirus by human monoclonal antibodies. *Journal of virology* 2014, 88:7796-805.
- [47] Wang N, Rosen O, Wang L, Turner HL, Stevens LJ, Corbett KS, Bowman CA, Pallesen J, Shi W, Zhang Y, Leung K, Kirchdoerfer RN, Becker MM, Denison MR, Chappell JD, Ward AB, Graham BS, McLellan JS: Structural Definition of a Neutralization-Sensitive Epitope on the MERS-CoV S1-NTD. *Cell reports* 2019, 28:3395-405 e6.
- [48] Zhou H, Chen Y, Zhang S, Niu P, Qin K, Jia W, Huang B, Zhang S, Lan J, Zhang L, Tan W, Wang X: Structural definition of a neutralization epitope on the N-terminal domain of MERS-CoV spike glycoprotein. *Nature communications* 2019, 10:3068.
- [49] Walls AC, Tortorici MA, Bosch BJ, Frenz B, Rottier PJM, DiMaio F, Rey FA, Veerles D: Cryo-electron microscopy structure of a coronavirus spike glycoprotein trimer. *Nature* 2016, 531:114-7.
- [50] Yuan M, Wu NC, Zhu X, Lee CD, So RTY, Lv H, Mok CKP, Wilson IA: A highly conserved cryptic epitope in the receptor-binding domains of SARS-CoV-2 and SARS-CoV. *Science* 2020.

- [51] McLellan JS, Chen M, Kim A, Yang Y, Graham BS, Kwong PD: Structural basis of respiratory syncytial virus neutralization by motavizumab. *Nature structural & molecular biology* 2010, 17:248-50.
- [52] Sanders RW, Derking R, Cupo A, Julien JP, Yasmeen A, de Val N, Kim HJ, Blattner C, de la Pena AT, Korzun J, Golabek M, de Los Reyes K, Ketas TJ, van Gils MJ, King CR, Wilson IA, Ward AB, Klasse PJ, Moore JP: A next-generation cleaved, soluble HIV-1 Env trimer, BG505 SOSIP.664 gp140, expresses multiple epitopes for broadly neutralizing but not non-neutralizing antibodies. *PLoS pathogens* 2013, 9:e1003618.
- [53] McLellan JS, Chen M, Joyce MG, Sastry M, Stewart-Jones GB, Yang Y, Zhang B, Chen L, Srivatsan S, Zheng A, Zhou T, Graepel KW, Kumar A, Moin S, Boyington JC, Chuang GY, Soto C, Baxa U, Bakker AQ, Spits H, Beaumont T, Zheng Z, Xia N, Ko SY, Todd JP, Rao S, Graham BS, Kwong PD: Structure-based design of a fusion glycoprotein vaccine for respiratory syncytial virus. *Science* 2013, 342:592-8.
- [54] Xiong H, Wu Y, Cao J, Yang R, Ma J, Qiao X, Yao X, Zhang B, Zhang Y, Hou W, Shi Y, Xu J, Zhang L, Wang S, Fu B, Yang T, Ge S, Zhang J, Yuan Q, Huang B, Li Z, Zhang T, Xia N: Robust neutralization assay based on SARS-CoV-2 S-bearing vesicular stomatitis virus (VSV) pseudovirus and ACE2-overexpressed BHK21 cells. *bioRxiv* 2020, doi: <https://doi.org/10.1101/2020.04.08.026948>.
- [55] Wu Y, Li C, Xia S, Tian X, Wang Z, Kong Y, Gu C, Zhang R, Tu C, Xie Y, Lu L, Jiang S, Ying T: Fully human single-domain antibodies against SARS-CoV-2. *bioRxiv* 2020, doi: <https://doi.org/10.1101/2020.03.30.015990>.
- [56] Ju B, Zhang Q, Ge X, Wang R, Yu J, Shan S, Zhou B, Song S, Tang X, Yu J, Ge J, Lan J, Yuan J, Wang H, Zhao J, Zhang S, Wang Y, Shi X, Liu L, Wang X, Zhang Z, Zhang L: Potent human neutralizing antibodies elicited by SARS-CoV-2 infection. *bioRxiv* 2020, doi: <https://doi.org/10.1101/2020.03.21.990770>.
- [57] Wang C, Li W, Drabek D, Okba NMA, van Haperen R, Osterhaus ADME, van Kuppeveld FJM, Haagmans BL, Grosveld F, Bosch B-J: A human monoclonal antibody blocking SARS-CoV-2 infection. *bioRxiv* 2020, doi: <https://doi.org/10.1101/2020.03.11.987958>.
- [58] Shan C, Yao Y-F, Yang X-L, Zhou Y-W, Wu J, Gao G, Peng Y, Yang L, Hu X, Xiong J, Jiang R-D, Zhang H-J, Gao X-X, Peng C, Min J, Chen Y, Si H-R, Zhou P, Wang Y-Y, Wei H-P, Pang W, Hu Z-F, Lv L-B, Zheng Y-T, Shi Z-L, Yuan Z-M: Infection with Novel Coronavirus (SARS-CoV-2) Causes Pneumonia in the Rhesus Macaques. DOI:1021203/rs225200/v1 2020.
- [59] Bao L, Deng W, Huang B, Gao H, Liu J, Ren L, Wei Q, Yu P, Xu Y, Qi F, Qu Y, Li F, Lv Q, Wang W, Xue J, Gong S, Liu M, Wang G, Wang S, Song Z, Zhao L, Liu P, Zhao L, Ye F, Wang H, Zhou W, Zhu N, Zhen W, Yu H, Zhang X, Guo L, Chen L, Wang C, Wang Y, Wang X, Xiao Y, Sun Q, Liu H, Zhu F, Ma C, Yan L, Yang M, Han J, Xu W, Tan W, Peng X, Jin Q, Wu G, Qin C: The Pathogenicity of SARS-CoV-2 in hACE2 Transgenic Mice. *bioRxiv* 2020, doi: <https://doi.org/10.1101/2020.02.07.939389>.
- [60] Brandi N, Williamson FF, Benjamin Schwarz, Kimberly Meade-White, Danielle P. Porter, Jonathan Schulz, Neeltje van Doremalen, Ian Leighton, Claude Kwe Yinda, Lizzette Pérez-Pérez, Atsushi Okumura, Jamie Lovaglio, Patrick W. Hanley, Greg Saturday, Catharine M. Bosio, Sarah Anzick, Kent Barbian, Tomas Cihlar, Craig Martens, Dana P. Scott, View ORCID Profile Vincent J. Munster, Emmie de Wit: Clinical benefit of remdesivir in rhesus macaques infected with SARS-CoV-2. 2020.
- [61] Diamond MS, Pierson TC: Molecular Insight into Dengue Virus Pathogenesis and Its Implications for Disease Control. *Cell* 2015, 162:488-92.

- [62] Ngoni AE, Shrestha S: Immune Response to Dengue and Zika. *Annual review of immunology* 2018, 36:279-308.
- [63] Casadevall A, Pirofski LA: The convalescent sera option for containing COVID-19. *The Journal of clinical investigation* 2020, 130:1545-8.
- [64] Liu L, Wei Q, Lin Q, Fang J, Wang H, Kwok H, Tang H, Nishiura K, Peng J, Tan Z, Wu T, Cheung KW, Chan KH, Alvarez X, Qin C, Lackner A, Perlman S, Yuen KY, Chen Z: Anti-spike IgG causes severe acute lung injury by skewing macrophage responses during acute SARS-CoV infection. *JCI insight* 2019, 4.
- [65] Tetro JA: Is COVID-19 receiving ADE from other coronaviruses? *Microbes and infection* 2020, 22:72-3.
- [66] Walls AC, Xiong X, Park YJ, Tortorici MA, Snijder J, Quispe J, Cameroni E, Gopal R, Dai M, Lanzavecchia A, Zamboni M, Rey FA, Corti D, Veesler D: Unexpected Receptor Functional Mimicry Elucidates Activation of Coronavirus Fusion. *Cell* 2019, 176:1026-39 e15.
- [67] Wan Y, Shang J, Sun S, Tai W, Chen J, Geng Q, He L, Chen Y, Wu J, Shi Z, Zhou Y, Du L, Li F: Molecular Mechanism for Antibody-Dependent Enhancement of Coronavirus Entry. *Journal of virology* 2020, 94.
- [68] Neher RA, Dyrda R, Druelle V, Hodcroft EB, Albert J: Potential impact of seasonal forcing on a SARS-CoV-2 pandemic. *Swiss medical weekly* 2020, 150:w20224.
- [69] Guan WJ, Liang WH, Zhao Y, Liang HR, Chen ZS, Li YM, Liu XQ, Chen RC, Tang CL, Wang T, Ou CQ, Li L, Chen PY, Sang L, Wang W, Li JF, Li CC, Ou LM, Cheng B, Xiong S, Ni ZY, Xiang J, Hu Y, Liu L, Shan H, Lei CL, Peng YX, Wei L, Liu Y, Hu YH, Peng P, Wang JM, Liu JY, Chen Z, Li G, Zheng ZJ, Qiu SQ, Luo J, Ye CJ, Zhu SY, Cheng LL, Ye F, Li SY, Zheng JP, Zhang NF, Zhong NS, He JX, China Medical Treatment Expert Group for C: Comorbidity and its impact on 1590 patients with Covid-19 in China: A Nationwide Analysis. *The European respiratory journal* 2020.
- [70] Pan Y, Zhang D, Yang P, Poon LLM, Wang Q: Viral load of SARS-CoV-2 in clinical samples. *The Lancet Infectious diseases* 2020, 20:411-2.

Figure legend

Figure1. SARS-CoV-2 structures, life cycle and inhibition strategies of virus replication. (A) The virus particle structure of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The spherical particle of SARS-CoV-2 consists of four structural proteins, including the spike proteins (S), the membrane protein (M), the envelope protein (E) and the nucleocapsid (N). The S, M, and E proteins are incorporated in the virus membrane, and the N protein is inside the particle and associated with virus genomic RNA. (B) The single-stranded positive-sense RNA genome of SARS-CoV-2. The ORF1a, or the ORF1a and ORF1b together, encodes the large polyproteins pp1a and pp1ab. The two polyproteins are further cleaved into 16 non-structural proteins (nsp1-nsp16). The papain-like protease (PLpro), the 3C-like protease (3CLpro), the RNA-dependent RNA polymerase (RdRp), and the exonuclease (ExoN) are indicated. In addition to the non-structural and structural proteins, the genome of SARS-CoV-2 also encodes six accessory proteins. (C) The structure of spike protein. The spike protein is a trimer and each monomer comprises an N-terminal S1 subunit and a C-terminal S2 subunit. The S1 subunit is further divided into the N-terminal domain (NTD), the receptor-binding domain (RBD), subdomain 1 (SD1), and subdomain 2 (SD2). The S2 subunit is further divided into the fusion peptide (FP), the heptad repeat 1 (HR1) and heptad repeat 2 (HR2). During virus entry, S1 is responsible for receptor binding, and S2 is responsible for membrane fusion. Figure 1B-C were not drawn proportional to their genome sizes. (D) The life cycle of SARS-CoV-2 and inhibition of virus replication with different strategies. Upon binding to the cellular receptor angiotensin-converting enzyme 2 (ACE2), the spike protein is activated by protease cleavage, such as TMPRSS2. After membrane fusion, the genomic RNA is released into the cytoplasm. The ORF1a and ORF1ab are translated into pp1a and pp1ab, which are cleaved by PLpro and 3CLpro to produce non-structural proteins. Several nsps, including nsp7-10, nsp12, and nsp14, form the replication-transcription machinery to produce genomic and subgenomic RNAs. After the translation of structural proteins, the virion is assembled at the ER-Golgi intermediate compartment (ERGIC) and encapsidate the N protein and genomic RNA. The mature virion is released outside of cells by transportation through the vesicles. Virus replication at the steps of entry, proteolysis, and genome replication can be inhibited as indicated.



UNCORRECTED

Table 1. Strategies for the discovery and development of SARS-CoV-2 neutralizing antibodies.

Target	The spike protein of SARS-CoV-2			
Antibody discovery	mAb sources	B cell from patient <ul style="list-style-type: none"> • Memory B cell • Plasma B cell 	Antibody library <ul style="list-style-type: none"> • Phage library • Yeast library • Ribosome library • Others 	Animal immunization <ul style="list-style-type: none"> • Mice (wild type or humanized) • Rabbit • Monkey • Lamma • Others
	mAb binding characteristics	<ul style="list-style-type: none"> • ELISA binding • Affinity measurement • Cell based binding • Epitope binning /mapping • Others 		
	Functional assays	<ul style="list-style-type: none"> • Blocking of spike (or RBD) binding to ACE2 • Blocking of spike-mediated membrane fusion • Pseudovirus based neutralization assay • Live virus based neutralization assay • Others 		
	Animal models	<ul style="list-style-type: none"> • Human ACE2 transgenic mice • Rhesus monkey • Ferrets • Cats • Others 		
Antibody development	Design of lead antibody	<ul style="list-style-type: none"> • Antibody humanization (if necessary) • Affinity maturation (if necessary) • Antibody format (single mAb, mAb cocktail or bi-specific) • Consideration of Fc-mediated effector function • Consideration of ADE • Others 		
	Preclinical development	<ul style="list-style-type: none"> • Antibody CMC (Chemistry, Manufacture and Control) • PK / PD models • Toxicity profile • Others 		
Clinical studies		<ul style="list-style-type: none"> • Clinical safety • Monotherapy • Combination therapy <ul style="list-style-type: none"> Antibodies targeting different epitopes Antibody + anti-viral small molecules Antibody + immune modulatory drugs (e.g. anti-IL6) • Cohort selection <ul style="list-style-type: none"> Patients with severe or mild symptoms Prophylaxis for high-risk people (e.g. front-line medical workers) 		

mAb: monoclonal antibody; ELISA: enzyme-linked immunosorbent assay; ACE2: angiotensin-converting enzyme 2; Fc: crystallizable fragment; ADE: antibody-dependent enhancement; PK: Pharmacokinetics; PD: pharmacodynamics; IL-6: Interleukine 6.

UNCORRECTED MANUSCRIPT