


Journal Pre-proof

Early mobilization and physical exercise and in patients with COVID-19: a narrative literature review

Veronica Lourenço Wittmer, Flavia Marini Paro, Halina Duarte, Verena Kise Capellini, Marcela Cangussu Barbalho-Moulim



PII: S1744-3881(21)00063-3

DOI: <https://doi.org/10.1016/j.ctcp.2021.101364>

Reference: CTCP 101364

To appear in: *Complementary Therapies in Clinical Practice*

Received Date: 22 August 2020

Revised Date: 13 February 2021

Accepted Date: 10 March 2021

Please cite this article as: Wittmer VL, Paro FM, Duarte H, Capellini VK, Barbalho-Moulim MC, Early mobilization and physical exercise and in patients with COVID-19: a narrative literature review, *Complementary Therapies in Clinical Practice*, <https://doi.org/10.1016/j.ctcp.2021.101364>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Elsevier Ltd. All rights reserved.

Physical exercise and early mobilization in patients with COVID-19: a narrative literature review

Author names:

Veronica Lourenço Wittmer^a, email: ve_lourenco@yahoo.com.br,

Flavia Marini Paro^a, email: flavia.paro@ufes.br,

Halina Duarte^a, email: halinaduarte@yahoo.com

Verena Kise Capellini^b, email: verena.capellini@unifesp.br

Marcela Cangussu Barbalho-Moulím^a, e-mail: mbarbalhomoulim@gmail.com

^aDepartment of Integrated Education on Health, Center of Health Science, Federal University of Espírito Santo (UFES), Vitória, ES, Brazil

^bDepartment of Biosciences, Institute of Health and Society, Campus Baixada Santista, Federal University of São Paulo (UNIFESP), Santos, SP, Brazil.

Declaration of competing interest (COI): The authors declare no conflict of interest.

Corresponding author:

Veronica Lourenço Wittmer, Centro de Ciências da Saúde, Departamento de Educação Integrada em Saúde, Av. Maruípe, 1468, Vitória, ES, Brazil, CEP: 29043-900, phone: 55-27-98185-5956

Early mobilization and physical exercise and in patients with COVID-19: a narrative literature review

Abstract

Background: Currently, little is known about early mobilization and exercise in individuals with COVID-19.

Objective: To describe the indication and safety of early mobilization and exercises in mild to severe COVID-19 patients and to investigate the use of telerehabilitation to deliver exercise programs to these patients.

Methods: This narrative literature review was conducted performing a comprehensive search of databases.

Results: 32 articles met the established criteria and the main findings were summarized and described, including indication, contraindication and recommendation for early rehabilitation and exercises prescription.

Conclusions: The literature suggests that early mobilization and physical exercise are beneficial for individuals with COVID-19. However, much of what has been published is based on expert opinion due to a lack of randomized trials, which are needed.

Keywords: COVID-19; Exercise, Early ambulation; Rehabilitation; Telerehabilitation, Physical Therapy.

1. Introduction

The newly emerging human coronavirus named severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the etiologic agent of human coronavirus

disease 2019 (COVID-19) [1]. After the first outbreak in Wuhan, China, this virus spread very quickly through Hubei Province [2] and worldwide, causing the current pandemic [1]. The rapid spread of the disease caused an unprecedented increase in demand for health care services and hospitalization raising work overload and requiring worldwide urgent efforts to reorganize every aspect of care [3]. This unexpected increase in demand for specialized acute care caused a deep crisis in the health system [4].

Although most people present mild forms of COVID-19, nearly 14% of individuals present severe forms, requiring hospitalization and oxygen therapy, and 5% need admission to intensive care units [5]. Depending on the COVID-19 severity, patients may suffer several dysfunctions, such as impaired lung function, physical deconditioning, muscle weakness [5], acute cerebrovascular diseases, venous thromboembolism [6], acute cardiac injury [7, 8, 9] psychological disorders, and cognitive impairments [5, 8, 10]. In general, critical COVID-19 patients present long length of hospital stay [5, 11] and prolonged bed rest and immobilization inevitably lead to several negative consequences [12, 13]. In addition, COVID-19 patients admitted to critical care often need long supportive care and mechanical ventilation, which put them at high risk to develop post-intensive care syndrome [14]. Furthermore, relatively young patients who survive acute respiratory distress syndrome can present persistent exercise limitations, reduced physical quality of life, and functional disability 5 years after their critical illness, which increases costs and use of health care services [15]. For all these reasons, early mobilization and exercises should be performed to prevent, relieve and/or recover these impairments and limitations, even in the acute stage of COVID-19 [5,11,16].

Early mobilization comprises different kinds of movements, from passive to

resistive movements up to activities, which are started immediately after the clinical stabilization [17]. It is worth mention that the term “activity” is being used as the execution of a task or action by an individual, according to The International Classification of Functioning, Disability and Health definition [18]. Early mobilization requires interdisciplinary team cooperation [19] and can be performed by different health professionals, especially physiotherapists [11, 20, 21], nurses [20] and occupational therapists [21]. On the other hand, exercise is defined as a subcategory of physical activity that is planned, structured, repetitive, and purposive, in the sense that the improvement or maintenance of one or more components of physical fitness is the objective [22]. Most of times, the exercises are prescribed and supervised by physiotherapists and exercise professionals.

The early mobilization and exercises are common routine in several health conditions [8, 9, 10, 33], as well as in critically ill patients [11, 12], and its effects are well established [8-12, 21, 22, 33]. However, as COVID-19 is a new disease, little is known about how and when mild, moderate or severe COVID-19 patients should be submitted to early mobilization and exercises. Besides that, the crescent use of remote tools for health services during the pandemic scenario, in an attempt to reduce the virus spread [31-33], reveals the need to know whether and how telerehabilitation has been used in COVID-19 patients.

Thus, the primary aim of this narrative review was to describe the indication and safety of early mobilization and exercises in mild to severe COVID-19 patients. The secondary objective was to investigate the use of telerehabilitation to deliver exercise programs to these patients.

It is important to highlight that respiratory physiotherapy for acute COVID-19 patients and rehabilitation for chronic sequelae after COVID-19 will not be debated

here since these subjects are out of the scope of this review and already have been discussed elsewhere [11,34-43).

2. Materials and methods

This narrative review was done following the methodology described by Gasparian et al [44].

2.1. Search strategy

A comprehensive search was conducted on MEDLINE, SciELO, PEDro and Web of Science up to June 30th 2020 to retrieve meta-analyses, systematic reviews, randomized trials, guidelines, recommendations, state of the art, and other peer-reviewed studies investigating the relationship between COVID-19 and early mobilization or exercises. Key terms, including Medical Subject Headings (MeSH) terms, were identified for each construct and combined using the Boolean operators “OR” within the construct and “AND” between constructs. The MeSH terms used were “COVID-19”, “SARS-CoV-2”, “severe acute respiratory syndrome coronavirus 2”, “coronavirus”, “exercise”, “early ambulation”, “physical therapy modalities”, “rehabilitation”, and “telerehabilitation”. The other key terms used were: “2019-nCoV”, “coronavirus pneumonia”, “physical exercises”, “physiotherapy”, “physical therapy”, “mobilization”, and “early mobilization”. All key terms were searched in titles and abstracts. All key terms were written in English, and when the database had the idiom filter tool, the language was restricted to English, Portuguese and Spanish (Table 1). To identify eligible papers that may not have been retrieved by the search described above, reference lists of included articles were manually checked. The search was conducted independently by two authors, who identified and remove duplicates, screened studies by title/abstract, and then reviewed full text, considering the inclusion

and exclusion criteria by eligibility. Discrepancies were resolved via consensus with a third author.

Table 1 should be inserted here.

2.2. Eligibility criteria

The inclusion criteria were studies where early mobilization and/or exercises were discussed for patients diagnosed with COVID-19.

The exclusion criteria were papers regarding exclusively the pediatric population, exclusively the respiratory exercise, and those that approach exclusively the rehabilitation after COVID-19.

3. Results

A total of 567 studies were identified in the searches. After the exclusion of 111 duplicate studies, 456 articles were screened by title and abstract, and the remained 85 articles were full text reviewed. So, 32 studies were included in this narrative review (Figure 1).

Figure 1 should be inserted here.

Among the 32 included papers, there are: thirteen opinion/perspective/communication/special/commentary/correspondence/experience/document reports, four letters to the editor, four types of reviews, four guidelines, three position papers, three editorials, and one randomized controlled trial (RCT) protocol (Table 2). It is necessary to highlight that there is no RCT investigating early mobilization and exercises in COVID-19 patients. Therefore, this narrative review is mainly based on observational studies in COVID-19, and on opinions of the scientific community about how to deal with early mobilization and exercises in COVID-19 patients, with these opinions arising from literature findings for other health conditions.

Table 2 should be inserted here.

The main findings of this review were grouped in the following topics: 3.1) COVID-19 manifestations and the need for early mobilization and exercises; 3.2) Early mobilization and exercises recommendations for mild to moderate COVID-19 patients; 3.3) Early mobilization and exercises recommendations for severe to critical COVID-19 patients; and 3.4) Telerehabilitation for COVID-19 patients.

3.1. COVID-19 manifestations and the need for early mobilization and exercises

Taking into account the multisystemic repercussions of COVID-19 [5,35,41,45-47] (Figure 2), which are common to other disease conditions, it is easy to understand the need of early mobilization and exercises to prevent, relieve and/or recover these impairments in body structures and functions, that ultimately will lead to activity limitations. And this need has been pointed out by recent guidelines [11,16,42,48,49] recommending early mobilization and exercises for COVID-19 patients to prevent and/or minimize these complications and to rehabilitate them. Besides the guidelines [11,16,42,48,49], all the other studies included in this review did also mention the importance of exercises for COVID-19 patients [35,40,46,50-52].

Figure 2 should be inserted here.

Although it is not yet possible to predict how many impairments will remain in post-acute and chronic phases, probably some patients will experience consequences during months or even longer, as observed on post intensive care syndrome cases [45]. Therefore, researchers, considering previous studies with critically ill patients, suggested that, in COVID-19 population, early mobilization and exercise can prevent muscle atrophy, venous thrombosis, stiffness of joints and soft tissue [49], prevent and reduce polyneuromyopathy, improve quality of life, reduce time and mortality during

hospitalization, contribute to the optimization of cognitive, respiratory, neuromuscular and osteoarticular function, shorten the duration of intensive care unit (ICU) stay and its clinical and functional sequelae [47], prevent and treat several adverse effects of sedation and bed rest [11,16,53,54]. Furthermore, the mechanical thromboprophylaxis (graduated compression stockings or intermittent pneumatic compression) additionally to pharmacologic thromboprophylaxis or as an option when pharmacological treatment is contraindicated was recommended for COVID-19 patients by the German Society of Angiology [55].

However, early mobilization and exercises prescription for COVID-19 patients should involve careful consideration of clinical conditions [11] and prescription parameters depend on the disease severity. As there are small variations regarding severity classification of COVID-19 [11, 16, 35, 40], this review adopted the World Health Organization (WHO)'s classification and definition of COVID-19 severity for adults [1] (Table 3) and, its next topics will report the indication (exercise type, intensity, frequency and duration) and safety (parameters to be monitored, exercise initiation and interruption criteria and contraindications), of early mobilization and exercises according to WHO COVID-19 severity classification.

Table 3 should be inserted here.

Still on exercise intensity, it is worth mention that moderate-intensity continuous training improves immune function biomarkers [56] and may optimize the functional integrity of the immune system to prevent or attenuate the severity of infection, especially among vulnerable populations with immune-compromised conditions [57]. Otherwise, high-intensity prolonged exercise leads to immunosuppression [58].

3.2 Early mobilization and exercises recommendations for mild to moderate COVID-19 patients

It is important to note that most of mild COVID-19 patients are not hospitalized [48,49], and then, they should be advised to stay home and balance rest and physical activity, paying attention to their symptoms. A home exercise program may be instituted, and an individualized approach is recommended specially in cases of immobility, neurologic, lung and cardiac diseases, and other complications.

For hospitalized COVID-19 patients, the modality, timing, and intensity of intervention for patients with COVID-19 must be tailored according to the individual patient's needs, especially for elderly, obese patients, and those with comorbidities [16]. Considering the individual's overall condition, the exercises intensity can be lowered to match the patients with reduced muscle strength [49]. Since some mild and moderate COVID-19 patients may rapidly progress to severe acute respiratory syndrome (SARS), it is recommended that interventions do not cause a further burden on breathing work, increasing the risk of respiratory distress [16].

The indication (exercise type, intensity, frequency and duration) and safety (parameters to be monitored, exercise initiation and interruption criteria and contraindications) of early mobilization and exercises in mild to moderate COVID-19 patients are summarized in Table 4.

Table 4 should be inserted here.

As can be seen in table 4, exercise intensity for mild COVID-19 patients should be limited to moderate. In fact, most of the studies recommended low-intensity exercise for these patients [16, 35, 51], represented by Borg category ratio (CR) $10 < 3$ or MET values < 3 . A single study advised low to moderate intensities (Borg CR 10 score ≤ 3) to mild COVID-19 patients [40], since the value 3 in the Borg CR 10 scale

corresponds to moderate intensity.

Regarding patients with moderate disease, some studies proposed low-intensity exercise (Borg CR 10 scale < 3 or MET values < 3) [16, 40], and other studies did not specify the intensity. However, the exercise intensity for moderate COVID-19 patients should not be high, since around 3% to 5% of moderately ill patients develop severe or critical disease after 7 to 14 days of infection; therefore, the aim of the exercise is to maintain the physical status [40].

Concerning complementary therapies, it was observed that an exercise program including elements of Health Qigong, exercises of pulmonary rehabilitation and proprioceptive neuromuscular facilitation (PNF), performed in 30-min session, daily, during 14 days of hospitalization of mild to moderate COVID-19 patients, improves functional activity. The authors of this pilot study reported that a standardized clinical research will be carried out [52]. Other authors also indicated Chinese traditional exercises (Tai chi chuan, Ba-duan-jin) for mild to moderate COVID-19 patients [35, 40].

3.3 Early mobilization and exercises recommendations for severe to critical COVID-19 patients

According to Li (2020) [35], there are still many doubts about rehabilitation of severe to critical COVID-19 patients; however, some consensus already have been achieved, such as the adoption of breath training at prone and/or semi recumbent bed position, moderate head elevation, limb mobilization, bed and bedside sitting and standing, as well as bedside walking when possible. Trying to better understand what “when possible” means, Kurtais et al [48] recommended that only passive range of motion exercises and secretion management should be done in COVID-19 patients presenting acute respiratory distress syndrome (ARDS). However, Righetti et al [51]

strongly recommended 12 to 16 hours/day of prone ventilation for severe ARDS patients and related that this maneuver can be repeated when the PaO₂/FiO₂ ratio of 150 mmHg is observed after 6 hours in the supine position, and must be interrupted if the patient presents hemodynamic instability. According to Lazzeri et al [42], as soon as clinical stability is reached, decubitus change, semi-prone, prone, semi-sitting and sitting position are recommended for severe patients spontaneously breathing or using non-invasive ventilatory support, and Kurtais et al [48] postulated that active exercises and electrical muscle stimulation (EMS) should be considered for those recovering from ARDS.

EMS applied in the early acute phase of severe COVID-19 patients has been indicated to counter the ICU acquired weakness [59]. Among the various EMS modes, belt-type is suggested as more effective for critical care, because it can induce whole lower extremity exercise through whole muscle contraction between wrapped belts, and 50 min bouts, three bouts per day have presented good results in ventilated COVID-19 patients [59]. Righetti et al [51] recommended that the EMS be associated with early verticalization and exercises in severe COVID-19 patients.

Early mobilization and exercises, including positioning with 45° head up in bed [60], regular bed decubitus changes [60], passive movement for all joints [16, 46, 60], frequent posture changes [16,46] with gradual increase of antigravity position in order to achieve semi-sitting position [46], and/or neuromuscular electrical stimulation [16], are also recommended for unconscious or sedated critical COVID-19 patients. These therapeutic approaches have to be choose based on patient's clinical condition [46], and are important to prevent skin lesions, immobilization sequelae [42] and disability [16]. The rehabilitation team should discuss the possibility to start an active mobilization program as soon as patient sedation is reduced, to avoid critical illness myopathy and

physical disability [42]. However, the active mobilization is not recommended for patients with clinical instability [42]. In these cases, the training of the skeletal muscle is contraindicated as it can impose an additional stress to the respiratory system, exposing the patients to an increased risk of respiratory distress [16].

Moreover, it already was demonstrated in severe COVID-19 patients that management of posture, passive and active mobilization, associated with respiratory and cognitive rehabilitation, depending on the clinical status, reduced the mean length of hospitalization and improved cognitive function in elderly who initially presented abnormal Mini-Mental State Examination or Montreal Cognitive Assessment scores [61].

Regarding safety, besides the contraindications and interruption criteria already described for mild to moderate COVID-19 patients (Table 4), severe to critical COVID-19 patients should also have the early mobilization and/or exercises stopped when disconnection of any device (intravenous access, endotracheal tube, feeding tube, etc.), irritability, heart palpitations, loss of consciousness, and falls appear [40].

3.4) Telerehabilitation for COVID-19 patients

An Australian guideline [11], that outlines recommendations for physiotherapy management for COVID-19 in the acute hospital setting, recommends the use of virtual tools to do an initial screening without direct contact with the patient whenever possible, calling the patient's isolation room telephone and conducting a subjective assessment and advising on patient mobility.

An editorial published [53] about physiotherapy in patients with COVID-19 also describe that, in collaborating with patients, it is possible to implement the execution of simple and low-intensity movements of upper and/or lower limbs in the sub-acute

settings. In addition, a guideline for acute and subacute rehabilitation of COVID-19 patients [48], recommends that the choice of printed and visual materials during telerehabilitation should depend on the patient's cognitive status.

A systematic review [41], cited the use of telerehabilitation to a patient with Charcot-Marie-Tooth disease who developed COVID-19 without major respiratory symptoms, and described one case series study, that applies a telerehabilitation system to deliver exercise to individuals who could not be assisted with the traditional face-to-face approach due to contagiousness concerns.

Most of the aforementioned studies [11, 41, 48, 61] corroborate that the telerehabilitation system is an important alternative to exercise with selected patients, without increasing the exposure and risk of contamination of professionals and patients.

A letter to the editor [62] described the experience obtained with the use of telerehabilitation for 4 patients hospitalized with Covid-19 through a tablet connected to the Internet via Wi-Fi and a pulse oximeter (used by the patients) connected to the tablet via Bluetooth. A physical therapist guided each individual in a 20-min exercise program, which consisted of stretching, muscle strengthening, and balance exercises directed by a video program with real-time instructions. On a scale of 0 to 10, the participant graduated their overall satisfaction between 8 and 10, and when they were asked "whether they felt it meaningful for their health to participate in this program", and "whether they would recommend this exercise to others" the answers ranged from 7 to 10 and from 8 to 10, respectively.

Lee et al.[63] reported that at the time their manuscript was wrote, as part of the Singapore government's containment effort, all individuals with COVID-19 were admitted into hospitals, although most of them were asymptomatic or have mild symptoms. So, exercise videos were provided to all patients in the general medical

wards to ensure that they remain active during their stay in the isolation facility [63]. The use of telerehabilitation in hospitalized patients with COVID-19 was also described for Falvey et al [64], and has emerged as an important strategy during this pandemic.

4. Discussion

Since Covid-19 is a new disease, there is still no robust scientific evidence to define the different aspects of early mobilization and physical exercise in affected patients. However, through our search we were able to verify, based on previous knowledge of the physiological effects of exercise, as well as on studies that investigated the safety and efficacy in patients with clinical situations similar to Covid-19, that exercise can probably be beneficial throughout pandemic context.

Although in our results we restricted our search to articles that address the effects of early mobilization and exercise only in Covid-19, we noticed during the reading of the articles presented, that many of them [11, 40, 41, 42, 46, 48] elaborated their recommendations based on studies carried out in other diseases, and only a few studies and experience reports [52, 59, 61, 63] cited effects and recommendations of exercise based on experience specifically with patients affected by Covid-19. The beneficial effects of exercise and mobilization in patients with other clinical conditions are well established in the literature [23, 29, 30]. However, mobilization is not recommended for patients with clinical instability [42], since it may not be well tolerated and cause rapid desaturation [45].

According to the guidelines of the American Association for Cardiovascular and Pulmonary Rehabilitation, the most challenging aspect of exercise prescription for pulmonary patients is the estimate of the appropriate intensity for each person to ensure that exercise is not too intense to cause adverse physiological effects, but it is enough to

promote beneficial effects [26]. Therefore, in the case of a new disease, about which there is a lack of scientific evidence, clinical decision making on the ideal intensity of exercise is even a greater challenge.

The need for exercise to rehabilitate patients with pneumonia caused by other etiological agents has been consolidated by some studies [23, 24]. Therefore, due the lack of RCTs about exercise on COVID-19, these studies about pneumonia have been considered for recommendations on COVID-19. Our results showed that most guidelines found [16, 35, 40, 51] recommends performing low intensity exercises to mild patients.

Regarding patients with severe and critical Covid-19 disease, our results also demonstrated that there are no RCTs that define which types of exercise and / or mobilization are safe and beneficial. However, all the articles selected in our search that addressed different forms of early mobilization [16, 35, 42, 48, 46, 51, 60, 61], highlighted that, since the safety criteria are respected, all Covid-19 patients should be mobilized precociously. While the patient is sedated or unconscious, passive movement of the patient [16, 42, 48]) and postural exchanges [16, 42, 60] are recommended, however, once the patient 'sedation is reduced, it is important to start the active mobilization program, to avoid physical disability and critical illness myopathy [42].

A systematic review demonstrated that early progressive mobilization in critically ill adults at Intensive Care Unit (ICU) is feasible and safe, improves functional ability and reduces ICU and hospital length of stay [30]. In addition, a RCT showed a reduction in delirium and duration of mechanical ventilation in critically ill patients who underwent early mobilization [65].

Since many patients with severe COVID-19 progress to mechanical ventilation and remain in hospital for long periods, they are likely to develop ICU-acquired

weakness. In this context, early mobilization can help to improve functional, cognitive, and respiratory conditions, and contribute to the patient's early discharge from the hospital [60].

Some studies [50, 56, 57] included in this review, evidenced that exercise can have positive enhancing effects on immune system and may be helpful for healthy asymptomatic people and, therefore, its practice is suitable during this viral respiratory tract epidemic [66]. Among the immunological effects of moderate exercise that have already been described, we can mention: increases neutrophil and natural killer (NK) cell counts [67] and augments stress hormones levels, which lead to reduction in excessive inflammation [67].

RCT studies are needed to investigate details about effect of exercise in patients with COVID-19, since, due to the unprecedented number of patients requiring intensive care, most publications addressing physical therapy and COVID-19 emphasize respiratory physical therapy. In the same way, there are several studies about exercise in post-discharge phase of COVID-19, since rehabilitation has been considered an urgent demand in this phase. The single RCT found during the screening process, about exercise in COVID-19 was conducted with patients after recovered from COVID-19 [36], however, it was excluded because it was an exclusion criterion. Our research also found a published protocol for a future RCT about the exercise in patients affected by COVID-19 confined to their homes, which will contribute to increase the knowledge about exercise in this population [68].

The restrictions imposed by the pandemic, can be a barrier to regular exercise and aggravate sedentary behaviors [69]. Special attention is deserved for the elderly population group, since exercise in older people positively affects and prevents frailty, sarcopenia, risk of falls, self-esteem and cognitive impairment [70].

Due to the aforementioned restrictions, typical rehabilitation programs need to be adapted [31, 32, 61, 71] and, probably for this reason, our results indicated a breakthrough in the use of virtual tools for exercise program delivery to patients COVID-19 [11, 63]. A RCT published before pandemic showed that telerehabilitation improved adherence to yoga in cardiac rehabilitation [72]. The use of telehealth technology is already a reality in many health services, with prospects for rapid expansion in the coming years, and development of different types of smart equipment, robots, wearable equipment for vital sign monitoring, among others, which are likely to perform a key role for prevention, treatment, and rehabilitation [35]. Telerehabilitation is an option for maintaining the mobility of the mild patients without increasing professional exposure, by mean of technologies to educate the patients on a safe exercise program and to encourage them to continue the exercise program independently [73]. For hospitalized and non-hospitalized stable patients in isolation, rehabilitation programs can be eventually conducted remotely by tele health system (educational videos, tele-consultation, webcams etc., with disinfectable tools) [73].

In conclusion, this review highlights the need for early mobilization and exercise to hospitalized patients with COVID-19 to prevent, reduce and rehabilitate the consequences of the disease and of the post-intensive care syndrome. Recommendation about early mobilization and exercise was described for patients with the mild, moderate, severe, or critical disease. Exercise prescription for these patients should be done with caution, observing each stage of the disease and the clinical condition, to ensure that exercise is not too intense to cause adverse physiological effects, but it is enough to promote beneficial effects. Although the available exercise recommendation is based on specialists' consensus and studies with low evidence level, the exposed data may help to guide the clinical practice of physical therapists, and other health

professionals until randomized controlled trials are published about exercise in COVID-19. Due to the occupational risk of contamination by the virus, telerehabilitation emerged as a valuable tool, especially in mild and stable COVID-19 patients.

References

- [1] World Health Organisation (WHO) Clinical management of COVID-19: interim guidance. 27 May 2020 Available at: <https://apps.who.int/iris/handle/10665/332196>
- [2] Novel coronavirus pneumonia emergency response epidemiology team. Vital surveillances: the epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) - China, 2020. China CDC Weekly. Accessed February 20, 2020. Available at: <http://weekly.chinacdc.cn/en/article/id/e53946e2-c6c4-41e9-9a9b-fea8db1a8f51>.
- [3] Willan J, King AJ, Jeffery K, Bienz N. Challenges for NHS hospitals during covid-19 epidemic. *BMJ*. 2020;368:m1117. DOI:10.1136/bmj.m1117.
- [4] Blumenthal D, Fowler EJ, Abrams M, Collins SR. Covid-19 - Implications for the Health Care System. *N Engl J Med*. 2020;383(15):1483-1488. doi: 10.1056/NEJMs2021088.
- [5] Pan American Health Organization (PAHO). Rehabilitation considerations during the COVID-19 outbreak. 2020. Available at: <https://iris.paho.org/handle/10665.2/52035>
- [6] Klok FA, Kruip MJHA, van der Meer NJM, Arbous MS, Gommers DAMPJ, Kant KM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res*. 2020;191:145-147. doi: 10.1016/j.thromres.2020.04.013.
- [7] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506. doi: 10.1016/S0140-6736(20)30183-5.

- [8] Clerkin KJ, Fried JA, Raikhelkar J, Sayer G, Griffin JM, Masoumi A, et al. COVID-19 and Cardiovascular Disease. *Circulation*. 2020;141(20):1648-1655. doi:10.1161/CIRCULATIONAHA.120.046941.
- [9] Zheng YY, Ma YT, Zhang JY, Xie X. COVID-19 and the cardiovascular system. *Nat Rev Cardiol*. 2020;17(5):259-260.
- [10] Carod-Artal FJ. Neurological complications of coronavirus and COVID-19. Complicaciones neurológicas por coronavirus y COVID-19. *Rev Neurol*. 2020;70(9):311-322.
- [11] Thomas P, Baldwin C, Bissett B, Boden I, Gosselink R, Granger CL, et al. Physiotherapy management for COVID-19 in the acute hospital setting: clinical practice recommendations. *J Physiother*. 2020;66(2):73-82.
- [12] Dittmer DK, Teasell R. Complications of immobilization and bed rest. Part 1: Musculoskeletal and cardiovascular complications. *Can Fam Physician*. 1993;39:1428-1437.
- [13] Prescott HC, Angus DC. Enhancing recovery from sepsis: A review. *JAMA*. 2018;319(1):62-75.
- [14] Jaffri A, Jaffri UA. Post-Intensive care syndrome and COVID-19: crisis after a crisis?. *Heart Lung*. 2020;49(6):883-884. doi:10.1016/j.hrtlng.2020.06.006.
- [15] Herridge MS, Tansey CM, Matté A, Tomlinson G, Diaz-Granados N, Cooper A, et al. Functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med*. 2011;364(14):1293-1304.
- [16] Vitacca M, Carone M, Clini EM, Paneroni M, Lazzeri M, Lanza A, et al. Joint statement on the role respiratory rehabilitation in the COVID-19 crisis: The Italian position paper. *Respiration*. 2020;1-7. doi:10.1159/000508399.
- [17] Arias-Fernández P, Romero-Martin M, Gómez-Salgado J, Fernández-García D.

Rehabilitation and early mobilization in the critical patient: systematic review. *J Phys Ther Sci.* 2018;30(9):1193-1201. doi:10.1589/jpts.30.1193.

[18] Towards a common language for functioning, disability and health: ICF - The International Classification of Functioning, Disability and Health. Geneva: World Health Organization; 2002. Available at: <https://www.who.int/classifications/icf/icfbeginnersguide.pdf>

[19] Hodgson CL, Stiller K, Needham DM, Tipping CJ, Harrold M, Baldwin CE, et al. Expert consensus and recommendations on safety criteria for active mobilization of mechanically ventilated critically ill adults. *Crit Care.* 2014;18(658):1-9.

[20] Garzon-Serrano J, Ryan C, Waak K, Hirschberg R, Tully S, Bittner EA, et al. Early mobilization in critically ill patients: patients' mobilization level depends on health care provider's profession. *PM R.* 2011;3(4):307-13. doi: 10.1016/j.pmrj.2010.12.022.

[21] Hashem MD, Nelliott A, Needham DM. Early mobilization and rehabilitation in the ICU: moving back to the future. *Respiratory care.* 2016;61(7), 971-979.

[22] Global action plan on physical activity 2018–2030: more active people for a healthier world. Geneva: World Health Organization; 2018. Licence: CC BY-NC-SA 3.0 IGO).

[23] Larsen T, Lee A, Brooks D, Michieli S, Robson M, Veens J, et al. Effect of Early Mobility as a Physiotherapy Treatment for Pneumonia: A Systematic Review and Meta-Analysis. *Physiother Can.* 2019;71(1):82-89. doi: 10.3138/ptc.2017-51.ep.

[24] José A, Dal Corso S. Inpatient rehabilitation improves functional capacity, peripheral muscle strength and quality of life in patients with community-acquired pneumonia: a randomised trial. *J Physiother.* 2016;62(2):96-102. doi: 10.1016/j.jphys.2016.02.014.

[25] Korkmaz C, Demirbas S, Vatansev H, Yildirim E, Teke T, Zamani A. Effects of

comprehensive and intensive pulmonary rehabilitation and nutritional support on quality of life and functional status in patients with chronic obstructive pulmonary disease. *J Int Med Res.* 2020;48(4):1-15.

[26] American Association of Cardiovascular & Pulmonary Rehabilitation (AACVPR). Guidelines for pulmonary rehabilitation programs. 6th edition. Champaign, IL, Human Kinetics; 2020

[27] Sawada Y, Sasabuchi Y, Nakahara Y, Matsui H, Fushimi K, Haga N, et al. Early rehabilitation and in-hospital mortality in intensive care patients with community-acquired pneumonia. *Am J Crit Care.* 2018; 27(2):97-103.

[28] Schujmann DS, Teixeira Gomes T, Lunardi AC, Zoccoler Lamano M, Fragoso A, Pimentel M, et al. Impact of a progressive mobility program on the functional status, respiratory, and muscular systems of ICU patients: A randomized and controlled trial. *Crit Care Med.* 2020;48(4):491-497.

[29] Momosaki R, Yasunaga H, Matsui H, Horiguchi H, Fushimi K, Abo M. Effect of early rehabilitation by physical therapists on in-hospital mortality after aspiration pneumonia in the elderly. *Arch Phys Med Rehabil.* 2015;96(2):205-209.

[30] Stiller K. Physiotherapy in intensive care: an updated systematic review. *Chest.* 2013;144:825-847.

[31] Chindhy S, Taub PR, Lavie CJ, Shen J. Current challenges in cardiac rehabilitation: strategies to overcome social factors and attendance barriers. *Expert Rev Cardiovasc Ther.* 2020;18(11):777-789. doi: 10.1080/14779072.2020.1816464.

[32] Ghram A, Briki W, Mansoor H, Al-Mohannadi AS, Lavie CJ, Chamari K. Home-based exercise can be beneficial for counteracting sedentary behavior and physical inactivity during the COVID-19 pandemic in older adults. *Postgrad Med.* 2020:1-12. doi: 10.1080/00325481.2020.1860394.

- [33] Babu AS, Arena R, Ozemek C, Lavie CJ. COVID-19: A time for alternate models in cardiac rehabilitation to take centre stage. *Can J Cardiol.* 2020;36(6):792-794. doi: 10.1016/j.cjca.2020.04.023.
- [34] Barker-Davies RM, O'Sullivan O, Senaratne KPP, Baker P, Cranley M, Dharm-Datta S, et al. The Stanford Hall consensus statement for post-COVID-19 rehabilitation. *Br J Sports Med.* 2020;54(16):949-959. doi: 10.1136/bjsports-2020-102596.
- [35] Li J. Rehabilitation management of patients with COVID-19. Lessons learned from the first experiences in China. *Eur J Phys Rehabil Med.* 2020. doi:10.23736/S1973-9087.20.06292-9.
- [36] Liu K, Zhang W, Yang Y, Zhang J, Li Y, Chen Y. Respiratory rehabilitation in elderly patients with COVID-19: A randomized controlled study. *Complement Ther Clin Pract.* 2020;39:1-4.
- [37] Rivera-Lillo G, Torres-Castro R, Fregonezi G, Vilaró J, Puppo H. Challenge for Rehabilitation After Hospitalization for COVID-19. *Arch Phys Med Rehabil.* 2020;101(8):1470-1471. doi: 10.1016/j.apmr.2020.04.013.
- [38] Wade DT. Rehabilitation after COVID-19: an evidence-based approach. *J Clin Med.* 2020;20(4):359-365.
- [39] Sheehy LM. Considerations for Postacute Rehabilitation for survivors of COVID-19. *JMIR Public Health Surveill.* 2020;6(2):e19462. doi: 10.2196/19462. PMID: 32369030; PMCID: PMC7212817.
- [40] Zhao H-M, Xie Y-X, Wang C. Recommendations for respiratory rehabilitation in adults with coronavirus disease 2019. *Chin Med J (Engl).* 2020;133(13):1595-1602.
- [41] Ceravolo MG, Arienti C, de Sire A, Andrenelli E, Negrini F, Lazzarini SG, et al. Rehabilitation and COVID-19: the Cochrane Rehabilitation 2020 rapid living systematic review. *Eur J Phys Rehabil Med.* 2020;56(5):642-651. doi: 10.23736/S1973-

9087.20.06501-6.

[42] Lazzeri M, Lanza A, Bellini R, Bellofiore A, Cecchetto S, Colombo A, et al. Respiratory physiotherapy in patients with COVID-19 infection in acute setting: a Position Paper of the Italian Association of Respiratory Physiotherapists (ARIR). *Monaldi Arch Chest Dis.* 2020;90:164-168.

[43] Severin R, Arena R, Lavie CJ, Bond S, Phillips SA. Respiratory Muscle Performance Screening for Infectious Disease Management Following COVID-19: A Highly Pressurized Situation. *Am J Med.* 2020;133(9):1025-1032. doi:10.1016/j.amjmed.2020.04.003

[44] Gasparyan AY, Ayvazyan L, Blackmore H, Kitas GD. Writing a narrative biomedical review: considerations for authors, peer reviewers, and editors. *Rheumatol Int.* 2011;31(11):1409-17. doi: 10.1007/s00296-011-1999-3.

[45] Kiekens C, Boldrini P, Andreoli A, Avesani R, Gamna F, Grandi M, et al. Rehabilitation and respiratory management in the acute and early post-acute phase. "Instant paper from the field" on rehabilitation answers to the COVID-19 emergency. *Eur J Phys Rehabil Med.* 2020;56(3):323-326.

[46] Brugliera L, Spina A, Castellazzi P, Cimino P, Tettamanti A, Houdayer E, et al. Rehabilitation of COVID-19 patients. *J Rehabil Med.* 2020;52(4):jrm00046. doi: 10.2340/16501977-2678. PMID: 32286674.

[47] Greve J M D, Brech GC, Quintana M, Soares A L de S, Alonso AC. Impacts of COVID-19 on the immune, neuromuscular, and musculoskeletal systems and rehabilitation. *Rev Bras Med Esporte.* 2020;26(4):285-288. <https://doi.org/10.1590/1517-869220202604esp002>.

[48] Kurtais Aytür Y, Köseoğlu BF, Özyemişçi Taşkiran Ö, Ordu-Gökkaya NK, Ünsal Delialioğlu S, et al. Pulmonary rehabilitation principles in SARS-COV-2 infection

(COVID-19): A guideline for the acute and subacute rehabilitation. *Turk J Phys Med Rehabil.* 2020;66(2):104-120. doi: 10.5606/tftrd.2020.6444.

[49] Rayegani SM, Raeissadat SA, Fakharian A, Babae M, Nezamabadi M, Boland Nazar NS, et al. Role of rehabilitation medicine in the COVID-19 pandemic: an Iranian consensus. *Eur J Phys Rehabil Med.* 2020. doi: 10.23736/S1973-9087.20.06366-2.

[50] Zhu Y, Wang Z, Zhou Y, Onoda K, Maruyama H, Hu C, et al. Summary of respiratory rehabilitation and physical therapy guidelines for patients with COVID-19 based on recommendations of World Confederation for Physical Therapy and National Association of Physical Therapy. *J Phys Ther Sci.* 2020;32(8):545-549. doi: 10.1589/jpts.32.545.

[51] Righetti RF, Onoue MA, Politi FVA, Teixeira DT, Souza PN, Kondo CS, et al. Physiotherapy Care of Patients with Coronavirus Disease 2019 (COVID-19) - A Brazilian Experience. *Clinics.* 2020;75:e2017. doi: 10.6061/clinics/2020/e2017.

[52] Chen JM, Wang ZY, Chen YJ, Ni J. The application of eight-segment pulmonary rehabilitation exercise in people with Coronavirus Disease 2019. *Front Physiol.* 2020;11:646. doi: 10.3389/fphys.2020.00646.

[53] Polastri M, Brini S, Ghetti A, Lama A. Recommendations from scientific/professional societies: An essential support for physiotherapy in patients with COVID-19. *Int J Ther Rehabil.* 2020;27(4). MA Healthcare Ltd. <https://doi.org/10.12968/ijtr.2020.0048>

[54] Polastri, M. Physiotherapy in hospitalised patients with COVID-19 disease: What we know so far. *Int J Ther Rehabil.* 2020;27(3). MA Healthcare Ltd. <https://doi.org/10.12968/ijtr.2020.0035>.

[55] Linnemann B, Bauersachs R, Grebe M, Klamroth R, Müller O, Schellong S, et al. Venous thromboembolism in patients with COVID-19 (SARS-CoV-2 infection) - a

position paper of the German Society of Angiology (DGA). *Vasa*. 2020;49:259-263.

[56] Rahmati-Ahmadabad S, Hosseini F. Exercise against SARS-CoV-2 (COVID-19): Does workout intensity matter? (A mini review of some indirect evidence related to obesity). *Obes Med*. 2020;19:100245. doi: 10.1016/j.obmed.2020.100245.

[57] Laddu DR, Lavie CJ, Phillips SA, Arena R. Physical activity for immunity protection: Inoculating populations with healthy living medicine in preparation for the next pandemic. *Prog Cardiovasc Dis*. 2020:S0033-0620(20)30078-5. doi: 10.1016/j.pcad.2020.04.006.

[58] Halabchi F, Ahmadinejad Z, Selk-Ghaffari M. COVID-19 Epidemic: Exercise or Not to Exercise; That is the Question! *Asian J Sports Med*. 2020;11(1):e102630, doi: 10.5812/asjasm.102630.

[59] Nakamura K, Nakano H, Naraba H, Mochizuki M, Hashimoto H. Early rehabilitation with dedicated use of belt-type electrical muscle stimulation for severe COVID-19 patients. *Crit Care*. 2020;24(1):342. doi: 10.1186/s13054-020-03080-5.

[60] Candan SA, Elibol N, Abdullahi A. Consideration of prevention and management of long-term consequences of post-acute respiratory distress syndrome in patients with COVID-19. *Physiother Theory Pract*. 2020;36(6):663-668. doi: 10.1080/09593985.2020.1766181.

[61] Iannaccone S, Castellazzi P, Tettamanti A, Houdayer E, Brugliera L, de Blasio F, et al. Role of Rehabilitation Department for Adult Individuals With COVID-19: The Experience of the San Raffaele Hospital of Milan. *Arch Phys Med Rehabil*. 2020;101(9):1656-1661. doi: 10.1016/j.apmr.2020.05.015.

[62] Mukaino M, Tatemoto T, Kumazawa N, Tanabe S, Katoh M, Saitoh E, et al. Staying Active in Isolation: Telerehabilitation for Individuals With the Severe Acute Respiratory Syndrome Coronavirus 2 Infection. *Am J Phys Med Rehabil*.

2020;99(6):478-479. doi: 10.1097/PHM.0000000000001441.

[63] Lee AJY, Chung CLH, Young BE, Ling LM, Ho BCH, Pua SH, et al. Clinical course and physiotherapy intervention in 9 patients with COVID-19. *Physiotherapy*. 2020;109:1-3. doi: 10.1016/j.physio.2020.06.002.

[64] Falvey JR, Ferrante LE. Flattening the disability curve: Rehabilitation and recovery after COVID-19 infection. *Heart Lung*. 2020;49(5):440-441. doi: 10.1016/j.hrtlng.2020.05.001.

[65] Schweickert WD, Pohlman MC, Pohlman AS, Nigos C, Pawlik AJ, Esbrook CL, et al. Early physical and occupational therapy in mechanically ventilated, critically ill patients: a randomised controlled trial. *Lancet*. 2009;373(9678):1874-1882.

[66] Thevarajan I, Nguyen THO, Koutsakos M, Druce J, Caly L, van de Sandt CE, et al. Breadth of concomitant immune responses prior to patient recovery: a case report of non-severe COVID-19. *Nat Med*. 2020;26(4):453-455.

[67] Martin SA, Pence BD, Woods JA. Exercise and Respiratory Tract Viral Infections: *Exerc Sport Sci Rev*. 2009;37(4):157-64.

[68] Gonzalez-Gerez JJ, Bernal-Utrera C, Anarte-Lazo E, Garcia-Vidal JA, Botella-Rico JM, Rodriguez-Blanco C. Therapeutic pulmonary telerehabilitation protocol for patients affected by COVID-19, confined to their homes: study protocol for a randomized controlled trial. *Trials*. 2020;21(1):588. doi: 10.1186/s13063-020-04494-w.

[69] Hall G, Laddu DR, Phillips SA, Lavie CJ, Arena R. A tale of two pandemics: How will COVID-19 and global trends in physical inactivity and sedentary behavior affect one another? *Prog Cardiovasc Dis*. 2020:S0033-0620(20)30077-3. doi: 10.1016/j.pcad.2020.04.005.

[70] Jiménez-Pavón D, Carbonell-Baeza A, Lavie CJ. Physical exercise as therapy to fight against the mental and physical consequences of COVID-19 quarantine: Special

focus in older people. *Prog Cardiovasc Dis.* 2020;63(3):386-388. doi: 10.1016/j.pcad.2020.03.009.

[71] Knudsen MV, Laustsen S, Petersen AK, Hjortdal VE, Angel S. Experience of cardiac tele-rehabilitation: analysis of patient narratives. *Disabil Rehabil.* 2019;1-8. doi: 10.1080/09638288.2019.1625450.

[72] Sarah S, Wolfgang MB, Claudia P. Effect of telerehabilitation on long-term adherence to yoga as an antihypertensive lifestyle intervention: Results of a randomized controlled trial. *Complement Ther Clin Pract.* 2019;35: 148-153. <https://doi.org/10.1016/j.ctcp.2019.02.001>

[73] Xie J, Tong Z, Guan X, Du B, Qiu H, Slutsky AS. Critical care crisis and some recommendations during the COVID-19 epidemic in China. *Intens Care Med.* 2020;46(5):837-840.

[74] Devlin JW, O'Neal HR Jr, Thomas C, Barnes Daly MA, Stollings JL, Janz DR, et al. Strategies to Optimize ICU Liberation (A to F) Bundle Performance in Critically Ill Adults with Coronavirus Disease 2019. *Crit Care Explor.* 2020;2(6):e0139. doi: 10.1097/CCE.0000000000000139.

[75] Simpson R, Robinson L. Rehabilitation After Critical Illness in People With COVID-19 Infection. *Am J Phys Med Rehabil.* 2020;99(6):470-474. doi: 10.1097/PHM.0000000000001443.

[76] Vitacca M, Lazzeri M, Guffanti E, Frigerio P, D'Abrosca F, Gianola S, et al. Italian suggestions for pulmonary rehabilitation in COVID-19 patients recovering from acute respiratory failure: results of a Delphi process. *Monaldi Arch Chest Dis.* 2020;90(2). doi: 10.4081/monaldi.2020.1444.




Figure 1. Flowchart detailing the systematic search, screening, eligibility, and inclusion procedure.




Figure 1. *Signs and symptoms* of coronavirus disease 2019 (*COVID-19*) and possible acute and chronic complications associated with *COVID-19* in different body systems.

Table 2. Characteristics of included studies (n = 34) and information about localization of papers findings in this narrative review.

(continue)

Author	Year	Country	Study design
Brugliera et al. [46]	2020	Italy	Letter to the editor (experience report)
Canan et al. [69]	2020	Turkey	Letter to the editor
Ceravolo et al. [41]	2020	Italy	Rapid living review
Chen et al. [52]	2020	China	Opinion article
Devlin et al. [74]	2020	United States	Review article
Falvey et al. [64]	2020	United States	Editorial
Gonzalez-Gerez et al. [68]	2020	Spain	Protocol
Greve et al. [47]	2020	Brazil	Special article (opinion of specialist)
Halabchi et al. [58]	2020	Iran	Editorial
Iannaccone et al. [61]	2020	Italy	Special communication
Kiekens et al. [54]	2020	Portuguese, Spanish	Special article
Kurtaiş et al. [48]	2020	Turkey	Guideline
Laddu et al. [57]	2020	United States	Commentary
Lazzeri et al. [42]	2020	Italy	Italian position paper
Lee et al. [63]	2020	China	Case series (short communication)
Li et al. [37]	2020	China	Special article
Linnemann et al. [55]	2020	Germany	Position paper
Mukaiho et al. [62]	2020	Japan	Letter to the editor
Nakamura et al. [59]	2020	Japan	Editorial
Pan American Health Organization [65]	2020	United States	PAHO Document
Polastri et al. [53]	2020	Italy	Editorial

Table 1. Search strategies used in each database.

Database	Search strategy
Pubmed	("COVID-19"[Title/Abstract] OR "SARS-CoV-2"[Title/Abstract] OR "severe acute respiratory syndrome 2019"[Title/Abstract] OR "coronavirus"[Title/Abstract] OR "2019-nCoV"[Title/Abstract] OR "coronavirus pneumonia"[Title/Abstract]) AND ("exercise"[Title/Abstract] OR "early ambulation"[Title/Abstract] OR "physical therapy modalities"[Title/Abstract] OR "rehabilitation"[Title/Abstract] OR "tele-rehabilitation"[Title/Abstract] OR "physical exercises"[Title/Abstract] OR "physiotherapy"[Title/Abstract] OR "physical therapy"[Title/Abstract] OR "early mobilization"[Title/Abstract] OR "early ambulation"[Title/Abstract]) Filters (Language: English OR Spanish OR Portuguese)
Web of Science	("COVID-19" OR "SARS-CoV-2" OR "severe acute respiratory syndrome 2019" OR "coronavirus" OR "2019-nCoV" OR "coronavirus pneumonia" OR "early ambulation" OR "physical therapy modalities" OR "rehabilitation" OR "physical exercises" OR "physiotherapy" OR "physical therapy" OR "early mobilization" OR "early ambulation")) Filters (Language: English OR Spanish OR Portuguese)
Scielo	("COVID-19 OR SARS-CoV-2 OR coronavirus OR "severe acute respiratory syndrome 2019" OR "2019-nCoV" OR "coronavirus pneumonia" OR "early ambulation" OR "physical therapy modalities" OR "rehabilitation" OR "physical exercises" OR "physiotherapy" OR "physical therapy" OR "early mobilization" OR "early ambulation"))). Limits: English, Portuguese and Spanish
PEDro	Search 05: COVID-19 AND exercise; Search 2: COVID-19 AND "early mobilization"; Search 3: COVID-19 AND physiotherapy; Search 4: COVID-19 AND "physical therapy"

Table 2. Characteristics of included studies (n = 34) and information about localization of papers findings in this narrative review.

(conclusion)

Author	Year	Country	Study design
Polastri et al. [54]	2020	Italy	Editorial
Rahmati-Ahmadabad & Hosseini [56]	2020	Iran	Correspondence
Rayegani et al. [49]	2020	Iran	Iranian consensus (letter to the editor)
Righetti et al. [51]	2020	Brazil	Experience report and review
Simpson & Robinson [75]	2020	Canada	Review
Thevarajan et al. [66]	2020	Australia	Correspondence
Thomas et al. [11]	2020	Australia	Guideline
Vitacca et al. [16]	2020	Italy	Italian position paper
Vitacca et al. [76]	2020	Italy	Consensus
Zhao H-M e at. [40]	2020	China	Guideline
Zhu et al. [50]	2020	China	Guideline

Table 3. WHO classification and definition of COVID-19 severity for adults [1].

Severity	Definition
Mild disease	Symptomatic patients meeting the criteria for diagnosis of COVID-19 without evidence of viral pneumonia or hypoxia.
Moderate disease (Pneumonia)	Adult with clinical signs of pneumonia, like fever, cough, dyspnea, tachypnea, however without signs of severe pneumonia, including $SpO_2 \geq 90\%$ (ambient air).
Severe disease (Severe pneumonia)	Adult with clinical signs of pneumonia, like fever, cough, dyspnea, tachypnea, with one of the following signs: respiratory rate > 30 breaths/min; severe respiratory distress; or $SpO_2 < 90\%$ (ambient air).
Critical disease	Onset: within 1 week of a known clinical insult (i.e. pneumonia) or new or worsening respiratory symptoms. Chest imaging showing bilateral opacities, not fully explained by volume overload, lobar or lung collapse, or nodules. Origin of pulmonary infiltrates: respiratory failure not fully explained by cardiac failure or fluid overload. (Assessment, e.g. echocardiography, to exclude hydrostatic cause of infiltrates/edema if no risk factor present).
(ARDS)	Classification of ARDS according to oxygenation impairment. Mild ARDS: $200 \text{ mmHg} < PaO_2/FiO_2 \leq 300 \text{ mmHg}$ (with PEEP or CPAP $\geq 5 \text{ cmH}_2\text{O}$); Moderate ARDS: $100 \text{ mmHg} < PaO_2/FiO_2 \leq 200 \text{ mmHg}$ (with PEEP $\geq 5 \text{ cmH}_2\text{O}$); Severe ARDS: $PaO_2/FiO_2 \leq 100 \text{ mmHg}$ (with PEEP $\geq 5 \text{ cmH}_2\text{O}$).
(Sepsis)	Adults with acute life-threatening organ dysfunction caused by a dysregulated host response to suspected or diagnosed infection.
(Septic shock)	Adults: persistent hypotension despite volume resuscitation, needs vasopressors to maintain MAP $\geq 65 \text{ mmHg}$ and serum lactate level $> 2 \text{ mmol/L}$.

WHO: World Health Organization; COVID-19: coronavirus disease 2019; SpO_2 : saturation of peripheral oxygen; ARDS: acute respiratory distress syndrome; FiO_2 : fraction of inspired oxygen; PEEP: positive end-expiratory pressure; CPAP: continuous positive airway pressure; MAP: mean arterial pressure.

Journal Pre-proof

Journal Pre-proof

Table 4. Recommendations about exercise for hospitalized patients with mild and moderate COVID-19.

Study	COVID-19 classification	Exercise type	Exercise intensity/ frequency/ duration criteria for interruption	Recommendation
Zhao et al.[34] 2020 China	Mild	Breathing exercises, Tai chi chuan, or square dancing.	Borg CR 10 scale dyspnea score ≤ 3 Twice a day, 15–45 min/session.	Start 1 h after meals; Fatigue should be absent on the next day (preferably).
Li [17] 2020 China	Mild Acute phase	Respiratory training, aerobic training, Chinese traditional exercises (Tai chi chuan, Ba-duan-jin), square dance.	Mild aerobic training.	
Kurtaiş et al. 2020 Turkey	Mild pneumonia	Active-assisted or active ROM exercises, achieving mobilization, muscle strengthening exercises, incentive spirometry and other devices for patients having sputum and productive cough, breathing techniques.		Started when the virulence decreases and when patient's condition is stabilized: decreased fever, reduced dyspnea, RR < 30 breaths/min, SpO ₂ > 90%.
Righetti et al.[36] 2020 Brazil	Mild Acute phase	Neuromuscular stimulation, therapeutic exercises, and early verticalization.	Light intensity exercises; the exercises can be for maintenance of a Borg CR 10 scale < 3.	To maintain minimal functional capacity. Start as soon as possible, as long as the patient presents suitable clinical conditions.
Zhu et al 2020 China	Mild/ Moderate Stable period	Respiratory training and RMT, mild exercise training, trying to walk, bedside bicycling, sitting and standing, resistance training, balance training; psychological support.	Exercises with intensity less than 3 Mets, maintaining a Borg score of 3–4 or visual analog scale (VAS) score 5–6. Progressive resistance training, three groups per day, 10 times per group, with a 1 repetition maximum of 50–70%. <u>Criteria for exercise interruption:</u> SpO ₂ decreased by 4 percentage points; or the patient had perspiration, nausea and vomiting, dizziness, blurred vision, etc.; severe sudden dyspnea; chest compression or pain; rapid heart rate or arrhythmia; unable to maintain balance.	<u>Initiation Criteria:</u> FiO ₂ $\leq 60\%$; SpO ₂ $\geq 93\%$; RR ≤ 30 breaths/min; PEEP ≤ 10 cmH ₂ O; SBP 90-180 mmHg; MAP 65-110 mmHg; HR 40-120 BPM; temperature < 38,5°. No new arrhythmias or myocardial ischemia; no new unstable deep vein thrombosis and pulmonary embolism; no aortic stenosis; no severe hepatorenal disease or new, progressive impairment of liver and kidney function. <u>Contraindications:</u> Temperature >38°; severe dyspnea; resting heart rate >120 BPM; X-ray manifestations: progression of thoracic infiltration >50% within 24–48 hours; SpO ₂ < 95%; BP < 90/60 mmHg or > 140/90 mmHg.

Table 4. Recommendations about exercise for hospitalized patients with mild and moderate COVID-19.

(Continuation)

Study	COVID-19 classification	Exercise type	Exercise intensity/ frequency/ duration criteria for interruption	Recommendation
Vitacca et al.[7] 2020 Italy	Mild/ Moderate Acute phase	Mobilization to get patient out of bed, posture changes/rotational therapy, therapeutic postures (early sitting/pronation), active limb exercises (also with devices), muscle reconditioning and strengthening, NES, RMT.	<u>Criteria for exercise withdraw</u> : high fever, worsening dyspnea, SpO ₂ < 93% or 4-point drop during exercise, chest tightness, belching, dizziness, headache, unclear vision, palpitations, sweating, inability to keep balance, increased need for O ₂ or NIV, radiological lesions' progression (> 50%) within 24-48h.	
Rayegani et al. 2020 Iran		Rolls on the bed, gets out of bed, sits on a chair, stands up, walks a few steps, stroll up and down the stairs, tai chi, preventive exercises for venous thrombosis, and Zheng's recumbent exercises. Active and passive joints range of motion exercises. The passive exercises are carried out if a patient has a reduced level of consciousness (assistive device for mobilization, ROM and stretching exercises, NES and compression stockings to prevent venous thrombosis). A full surface massage while using infrared light and TENS therapy to reduce myalgia.	The intensity of physical activity between 1-3 METs. Twice a day, at least 1 hour after eating, and between 15 and 45 minutes, as beyond that may cause fatigue in patients.	
Zhao et al.[34] 2020 China	Moderate	Breathing exercises, stepping, Tai chi chuan and exercises to prevent thrombosis.	1 < METs to < 3 METs (from rest to light) Twice a day, 15–45 min/session. Patients who are prone to fatigue or are physically weak should perform intermittent exercise. 3–5% of these patients develop severe disease after 7–14 d. So, intensity should not be high (exercise objective: maintain previous physical state) <u>Criteria for exercise interruption</u> : Borg > 3 (dyspnea), breathlessness, chest tightness, headache, blurred vision, palpitations, profuse sweating, dizziness, balance disorder, etc.	Start 1 h after meals <u>Exclusion criteria</u> : T > 38°C, initial consultation time ≤ 7 d, from disease onset to dyspnea ≤ 3 d, chest radiological scans > 50% progression within 24–48 h, SpO ₂ ≤ 95%, resting BP: <90/60 or >140/90 mmHg.

Table 4. Recommendations about exercise for hospitalized patients with mild and moderate COVID-19.

(Conclusion)

Study	COVID-19 classification	Exercise type	Exercise intensity/ frequency/ duration criteria for interruption	Recommendation
Thomas et al.[6] 2020 Australia		Range of motion (passive, active-assisted, active or resisted), muscle strength, bed mobility, sitting out of bed, sitting balance, sit to stand, walking, tilt table, standing hoists, upper/lower ergometry and exercise programs.	Mobilization and exercise prescription depends on the patient clinical status.	Direct physiotherapy interventions should only be considered when there are significant functional limitations, such as (risk of) ICU-acquired weakness, frailty, multiple comorbidities and advanced age.

BP: blood pressure; COVID-19: coronavirus disease 2019; CR: category ratio; h: hour(s); d: days; FiO₂: fraction of inspired oxygen; HR: heart rate; ICP: intracranial pressure; MRC: Medical Research Council; MAP: mean arterial pressure; MET: metabolic equivalent of task; NES: neuromuscular electrical stimulation; NIV: non-invasive ventilation; PEEP: positive end expiratory pressure; RASS: Richmond agitation sedation scale; RMT: respiratory muscle training; ROM: range of motion; RR: respiratory rate; SpO₂: peripheral capillary oxygen saturation; SBP: systolic blood pressure; T: temperature

Journal Pre-proof

Highlights

Mobilization plans must consider disease severity, contraindications and comorbidities.

Early mobilization should start as soon as COVID-19 patients have clinical stability.

Exercise prescription must consider clinical conditions and contraindications.

Most of the studies recommended low-intensity exercise for patients with mild disease.

Telerehabilitation should play a key role for patients in the recovery of COVID-19.