

Journal Pre-proof

Self-reported and clinically identified loss of smell and taste among persons tested for COVID-19 in Chennai, southern India, july–august 2020: A cross sectional study

Kathiresan Jeyashree, Mohankumar Raju, Manickam Ponnaiah, Sendhilkumar Muthappan, Amanda G.A. Rozario, Rose Raichel, W. Lydia Jeris, Raman R. Gangakhedkar, Manoj V. Murhekar

PII: S2213-3984(21)00022-1

DOI: <https://doi.org/10.1016/j.cegh.2021.100718>

Reference: CEGH 100718

To appear in: *Clinical Epidemiology and Global Health*

Received Date: 29 December 2020


Revised Date: 3 March 2021

Accepted Date: 8 March 2021

Please cite this article as: Jeyashree K, Raju M, Ponnaiah M, Muthappan S, Rozario AGA, Raichel R, Jeris WL, Gangakhedkar RR, Murhekar MV, Self-reported and clinically identified loss of smell and taste among persons tested for COVID-19 in Chennai, southern India, july–august 2020: A cross sectional study, *Clinical Epidemiology and Global Health* (2021), doi: <https://doi.org/10.1016/j.cegh.2021.100718>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier B.V. on behalf of INDIACLEN.



Self-reported and clinically identified loss of smell and taste among persons tested for COVID-19 in Chennai, Southern India, July-August 2020: A cross sectional study

Kathiresan Jeyashree^a, Mohankumar Raju^a, Manickam Ponnaiah^a, Sendhilkumar Muthappan^a, Amanda G. A Rozario^a, Rose Raichel^b, W. Lydia Jeris^b, Raman R Gangakhedkar^c, Manoj V Murhekar^{a*}

^a ICMR-National Institute of Epidemiology, Chennai, Tamil Nadu, India

^b Dr.Mehta's Hospitals, Chennai, Tamil Nadu, India

^c Division of Epidemiology & Communicable Diseases, Indian Council of Medical Research, New Delhi, India

* Corresponding author

Manoj V Murhekar, MD
ICMR-National Institute of Epidemiology,
R 127, TNHB, Ayapakkam,
Chennai, Tamil Nadu, 600 077, India
Email: mmurhekar@gmail.com

Running title: Loss of smell and taste in COVID-19

Number of words in manuscript: 2861

Total references: 43

Tables: 4

Figures: 1

1 Abstract

2 Background

3 Early detection of symptoms of loss of smell and taste lately added for Coronavirus disease
4 2019 (COVID-19) has the potential for improving pandemic response. In the Indian context,
5 we compared proportion experiencing new loss of smell or taste among COVID-19 positive
6 and negative individuals in Chennai city, Southern India.

7 Methods

8 We did an analytical cross-sectional study among aged 18-80 years undergoing testing at
9 COVID-19 sample collection centres. We ascertained loss of smell and taste using
10 standardised self-reporting and clinical examination procedures. We administered Sino Nasal
11 Outcome (SNOT 22) questionnaire for comprehensive understanding of these symptoms. We
12 compared proportion having symptoms between COVID-19 positive and negative persons.
13 We compared the two assessment methods to compute diagnostic validity indicators.

14 Results

15 Of the 277 participants, 169 (61%) were men and mean age of 40.7 years [SD=13.3]. 58
16 (21%) had COVID-19 and 12 (36%) of them were asymptomatic. Predominantly reported
17 symptoms were fever (30%), headache (18%) and cough (18%). Self-reported or clinically
18 identified new loss of smell or taste was higher among COVID-19 positive (n=13; 22%) than
19 negative persons (n=23; 11%) [p=0.02]. Sensitivity was higher for self-reported or clinically
20 identified loss of smell (17.2%) than that of loss of taste (6.9%). Negative predictive value for
21 loss of smell or taste, self-reported or clinically identified was 81%. Likelihood ratio of
22 positive test was 2.13.

23 Conclusion

24 Loss of smell or taste are predominantly reported by COVID-19 confirmed individuals.
25 Objective and subjective assessments of smell and taste may be required to identify those
26 requiring COVID-19 testing.

27 **Keywords:** Anosmia; Dysgeusia; Gustation; Olfaction; SNOT

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

Self-reported and clinically identified loss of smell and taste among persons tested for COVID-19 in Chennai, Southern India, July-August 2020: A cross sectional study

1. Introduction

As of October 2020, there have been over 40 million confirmed cases of Coronavirus disease 2019 (COVID-19), including around 1.1 million deaths worldwide.¹ India suffers one of the highest burden of cases with over 8 million cases recorded up to October 2020.² While still in pursuit of a vaccine and drug, the strategies employed to combat the pandemic are to test, treat and isolate, alongside community-based preventive measures emphasising social distancing, and hand hygiene among others. India's testing strategies have undergone revisions to suit the demands of the evolving pandemic and the country is currently testing for COVID-19 at the rate of around 19,000 tests per one million population.^{3,4} The current testing strategy advises testing those with symptoms of influenza-like-illness or severe acute respiratory illness and asymptomatic people with a history of direct, and high-risk contact with a confirmed case.⁵

A considerable proportion of the COVID-19 cases can be asymptomatic.⁶ Globally, the commonly reported symptoms among patients with COVID-19 include fever, cough, fatigue, sputum production, shortness of breath, myalgia, sore throat, headache, and others.⁷⁻¹⁰ Other gastrointestinal¹¹ and dermatological symptoms¹² have also been reported.

The United States Centers for Disease Control and Prevention (CDC) listed six new symptoms that were reported among patients with COVID-19 including new loss of taste (anosmia) or smell (ageusia), chills, and muscle ache.^{13,14} In a subsequent addition to this list

1 of symptoms, nausea or vomiting, runny nose, and diarrhea were also included.¹⁰ Anosmia,
2 with or without ageusia, has been reported commonly to manifest either early in the disease
3 process or in patients with mild or no constitutional symptoms.¹⁵⁻²¹ The prevalence of loss of
4 smell widely ranged from 10% to over 80% in European countries. Odor threshold detection
5 was more affected by COVID-19 compared to odor identification.²² The proportion reporting
6 taste disorders ranged up to 60%. The guidelines for diagnosis recommend chemosensory
7 assessment by subjective assessment and psychophysical assessment testing for one or a
8 combination of odor threshold, discrimination, and identification. Management of anosmia
9 includes safety counselling, olfactory training and adjuvant medication.²³

10 None of the listed symptoms are specific to COVID-19. Even seemingly specific symptoms
11 like loss of smell and taste are common with other neurotropic viral infections. Struyf et al in
12 their Cochrane review highlighted the low sensitivity and specificity of any one of the earlier
13 listed symptoms of COVID-19 and the need for data on potentially more specific symptoms
14 such as loss of sense of smell.²⁴ Menni et al have reported that loss of smell and taste is a
15 potential predictor of COVID-19 in addition to other, more established,
16 symptoms.²⁵ However, olfactory and gustatory dysfunction do not seem to hold prognostic
17 value at the time of initial diagnosis.²⁶

18 It is essential to understand the proportion of patients in India who present with these newly
19 listed symptoms, some of which are not otherwise routinely enquired or documented. This
20 will enable us to identify cases that would have earlier been missed and follow up with timely
21 isolation, and effective management.

22 In this context, we aimed to (1) describe the symptom profile of persons who underwent
23 testing for COVID-19 in selected sample collection centres in Chennai, Southern India (2)
24 compare the proportion reporting the new loss of smell or taste between those who tested

1 COVID-19 positive and COVID-19 negative. Additionally, we estimated diagnostic validity
2 indicators of loss of sense of smell and taste towards screening for COVID-19.

3 2. Materials and Methods

4 2.1. Study setting

5 Chennai, the fourth largest city in India, is situated in the southern state of Tamil Nadu and
6 home to a population of over 7 million. The city had been reporting an average of 1,200
7 COVID-19 cases per day and ranks third among the country's cities in terms of the COVID-
8 19 case burden. Nasopharyngeal swab is collected from all persons who require testing for
9 COVID-19 at sample collection centres which transport them to designated laboratories.
10 Persons testing positive for COVID-19 are screened for the severity of disease at screening
11 centres and based on the severity of the disease, recommended home isolation, admission to
12 COVID care centres or dedicated COVID hospitals.

13 2.2. Study design

14 We conducted an analytical cross-sectional study.

15 2.3. Study population

16

17 Adults aged between 18-80 years belonging to any gender, who visited COVID-19 testing
18 centres in Chennai city in Southern India

19 2.4. Sample size

20

21 We had to recruit 55 COVID-19 positives and 165 COVID-19 negative individuals. Our
22 assumptions were prevalence of self-reported loss of smell or taste of 25% among COVID-19
23 positive [based on case series (unpublished) of self-reporting through telephonic interviews
24 of COVID-19 individuals] and 5% among COVID-19 negative persons, alpha error of 5%,
25 1:3 allocation ratio, and non-response of 20%. We enrolled all eligible study participants

1 consecutively from two COVID-19 specimen collection centres until desired sample size was
2 achieved

3 2.5. Data collection

4
5 The data was collected from all eligible and consenting participants. The data was collected in
6 two parts. A face-to-face data collection of minimal identification details and clinical
7 examination were conducted at the sample collection centre for COVID -19. This was
8 followed by telephonic interviews on the same day.

9 After enquiring about history of new onset loss of smell and taste in the past 14 days,
10 participants were clinically evaluated for loss of smell and taste using standardised tests in the
11 testing booths set up with all necessary COVID-19 infection control measures including
12 personal protection equipment, social distancing, and proper disinfection of testing material.
13 We completed the clinical examination for loss of smell and taste ahead of sample collection
14 for COVID-19 testing at the centres. The standardisation procedures used for the preparation
15 of solutions and testing were based on De Jong's 'The Neurologic Examination' [Box 1].²⁷
16 For standardising the solutions to be used for smell and taste testing, serially increasing
17 concentrations of the solutions were tested on ten apparently healthy participants. A
18 concentration higher than the highest concentration at which the smell or taste was perceived
19 by all volunteers was chosen for the study participants.

20 Following this, trained interviewers telephonically contacted the participants to collect data
21 on a structured questionnaire (developed using Open Data Kit (ODK) application). In order to
22 limit reporting biases, the data was collected on the same day of the specimen collection, and
23 prior to the declaration of COVID-19 test results. The interviewers collected data on
24 demographic details, symptoms and other co-morbidities, and administered Sino-Nasal

1 Outcome (SNOT 22) questionnaire for comprehensive understanding of their symptoms. The
2 nasal and oropharyngeal samples were collected and tested by RT-PCR for COVID-19 at the
3 Government accredited laboratories. The results of this testing was obtained from the
4 laboratory to update the participant COVID-19 status.

5 2.6. Data analysis

6 Socio-demographic characteristics and symptom profile were summarized as frequencies and
7 percentages. Loss of smell/taste, self-reported or clinically identified, were statistically
8 (Fisher's exact test) compared between COVID positive and negative participants. We
9 computed positive predictive value (PPV), Negative Predictive value (NPV) and likelihood
10 ratio of a positive test (LRpos) to describe the diagnostic properties of the two methods of
11 ascertaining loss of smell and taste, namely self-reporting and clinical examination. We used
12 Epi-Info (Version 7.2) for analysis.

13 2.7. Human participant protection

14 The study was approved by the Institutional Human Ethics Committee of the ICMR-National
15 Institute of Epidemiology, Chennai. We obtained the verbal informed consent from all the
16 participants before conducting the study.

17 3. Results

18

19 3.1. Profile of study participants

20 Of the total 277 persons studied, 169 (61%) were men, 247 (93%) could read and write, and
21 20 (8%) reported living below the poverty line. The mean (SD) age of the study participants
22 was 40.7 (13.3) years (Tables 1 and 2). Overall, 139 (50%) persons tested reported no
23 symptoms at all. 58 (21%) persons tested positive for COVID-19. Among 58 who tested
24 positive for COVID-19, 36 (64%) had reported at least one symptom. Fever (30%) was the

1 most common symptom reported by them, while cough (14%) was the most common
2 symptom reported by those who tested COVID-19 negative (Figure 1). The SNOT 22 score
3 [Median; Interquartile range (IQR)] was not different between the two groups [18 (18-20) vs.
4 19; (18-21)].

5 **3.2. Frequency of Loss of smell/taste**

6 Overall, 21 (7.6%) (95% CI: 5.1 to 12.1) persons self-reported loss of smell or taste. Clinical
7 examination identified 18 (6.5%) (95% CI: 4.3 to 11.0) with loss of smell or taste. It was
8 found that in total, 36 (13%) (95% CI: 9.9 to 18.7) persons had loss of smell or taste by self-
9 reporting or clinical examination. Among persons with COVID-19, clinical examination
10 identified more number of people suffering from loss of smell compared to self-reporting. On
11 the contrary, among COVID- negative persons, self-reported loss of smell was higher than
12 what was identified clinically (Table 3).

13 New loss of smell or taste was reported by or identified by clinical examination among 13
14 (22.4%) (95% CI: 13.2 to 31.0) participants who tested COVID-19 positive and among 23
15 (10.5%) (95% CI: 7.3 to 16.4) COVID-19 negative persons ($p=0.016$). A person with loss of
16 smell or taste had 2.46 (1.16, 5.23) times higher odds for testing COVID-19 positive
17 compared to a person who did not experience these symptoms.

18 There was no difference in median time (days) to testing from first symptom, among those
19 self-reporting loss of smell and taste (4 days for both) compared to those identified to have
20 the symptoms by clinical examination (smell=4; taste= 5 days).

21 **3.3. Diagnostic validity: Self-reporting Vs Clinical examination**

22 We calculated the negative predictive value of loss of smell or taste, self-reported or
23 clinically identified to be 81% (Table 4). The probability that someone is likely to test

1 negative for COVID-19 if they were determined to not have loss of smell or taste by self-
2 reporting or clinical examination was over 80%.

3 A person is almost two times ($LR_{pos}=2.13$) more likely to report loss of smell or taste if s/he
4 tested positive for COVID-19 as compared to someone who tested negative (Table 4). The
5 sensitivity of self-reported or clinically identified loss of smell was 17.2% and that of loss of
6 taste was 6.9%.

7 4. Discussion

8 We did an analytical cross-sectional study to ascertain loss of smell or taste through self-
9 reporting as well as clinically examination among persons reporting for COVID-19 testing by
10 RT-PCR at sample collection centres in Chennai city, Southern India. We identified that a
11 significantly higher proportion of COVID-19 positive persons experienced loss of sense of
12 smell or taste compared to those tested negative for COVID-19.

13 Of the 58 COVID-19 positive patients in our study, 35.7% were asymptomatic at the time of
14 presentation for testing. This is comparable to what was reported in earlier in India, where
15 around 28% of the 40,814 COVID-19 cases were asymptomatic.²⁸ Systematic reviews also
16 report the pooled proportion of asymptomatic COVID-19 infections to be around 10 to 20 %,
17 slightly higher in children than other age groups.^{6,29,30} The difference in estimates across
18 studies could also be due to differences across countries in testing strategies and these
19 systematic reviews did not include any studies from India. Fever and cough were the most
20 commonly reported symptoms among those who tested COVID-19 positive as reported in
21 other studies.^{10,31}

22 4.1. Loss of smell and taste

1 We report that almost one-fifth of the COVID-19 affected persons in our study experienced
2 loss of smell or taste. Mullol et al. observe that the frequency of smell or taste dysfunction in
3 COVID-19 affected persons has shown a high variability from 5% to 98%, depending on the
4 methodology, country, and study.³² Our estimates fall within the range estimated by Larco et
5 al. who report that the frequency of anosmia in COVID-19 affected persons ranged between
6 22%-68% and frequency of taste from 20%-33%.³³

7 However, our estimates of the prevalence of loss of smell and taste are lower than that from
8 other systematic reviews. Samaranayake et al. in their meta-analysis including 11,054
9 COVID-19 patients from eight studies estimated that anosmia and dysgeusia were present in
10 74.9%, and 81.3% mild-to-severe cases of COVID-19 patients, respectively.³⁴ The pooled
11 proportions presenting with olfactory dysfunction and gustatory dysfunction based on data
12 from 24 studies on 8,438 patients were 41.0% and 38.2%, respectively.³⁵ The overall
13 prevalence of self-reported alteration of the sense of smell or taste was 31% and 67% in
14 severe and mild-to-moderate symptomatic patients, respectively.³⁶

15 We observed a significantly higher proportion of COVID-19 positive persons reporting loss
16 of sense of smell and taste compared to those who were COVID-19 negative. Boscolo et al.
17 followed up household contacts of COVID-19 patients for symptoms, and found a
18 significantly lower prevalence of smell or taste impairment (1.5%) in patients who tested
19 negative compared to those who tested positive for SARS-CoV-2 (63.0%).^{37,38}

20 Considering the reported possibility of false-negative results with COVID-19 RT-PCR
21 testing,³⁹ there is a possibility that some of our COVID negative study participants were
22 actually COVID positive thereby giving an overestimate of the proportion of COVID
23 negative persons experiencing the symptoms. This differential misclassification could have
24 diluted the difference between the two study groups namely COVID positive and COVID

1 negative persons. Thus, the actual difference in the proportion reporting these symptoms
2 between COVID-19 positive and negative persons could be wider than what we have
3 reported in our study.

4 In our study, the proportion of self-reporting loss of smell or taste was different from those
5 detected by clinical examination. Prevalence of olfactory dysfunction is significantly greater
6 using objective olfactory assessments compared to subjective measures.^{35,40} Lechien et al also
7 report that of 18 (38.3%) individuals who self-reported subjective partial or total loss of
8 smell, only three were anosmic.⁴¹ The use of subjective measures, while operationally more
9 feasible, may lead to underestimation of true prevalence.⁴²

10 **4.2. Diagnostic validity: Self-reporting Vs Clinical examination**

11 Struyf et al reported sensitivity of 0.22 for loss of smell and 0.20 for loss of taste and a
12 specificity of 0.95 and 0.96 respectively for the same symptoms.²⁴ Our estimates of
13 specificity are comparable with these, while those of sensitivity are much lower. We report
14 higher odds of being COVID-19 positive among those reporting loss of smell or taste
15 compared to those who didn't. While other studies support this finding, their point estimate of
16 the measure of association is higher. Those who reported loss of smell and taste had six-fold
17 higher odds of being COVID-19 positive; similarly, anosmia and ageusia were associated
18 with 10-fold higher odds of COVID-19 diagnosis.³³ In populations of patients who are
19 currently reporting olfactory dysfunction, there was a positive predictive value of 61% for a
20 positive COVID-19 result.³⁸ One reason could be the subjective nature of the symptoms
21 reported which require a certain level of awareness and general health literacy among the
22 population studied. The estimates available in the literature are thus far from Western
23 countries with a higher literacy rate and a higher overall socioeconomic status compared to
24 our study population.

1 **4.3. Implications of the findings**

2 Our findings have implications for an individual, the clinicians and the public health experts.
3 An individual suffering from these symptoms is required to self-isolate himself while also
4 seeking testing for COVID-19 and has to be prescribed an appropriate line of management
5 thereafter. The practicing clinician should be astute to refer persons reporting to him with
6 these symptoms at the out-patient clinic or as in-patients for testing for COVID-19. Public
7 health officials may incorporate these symptoms into their screening tool to identify the
8 population requiring COVID-19 testing.

9 **4.4. Strengths**

10 To the best of our knowledge, this is one of the few studies from India to document the
11 proportion of COVID-19 affected persons suffering from loss of smell and taste. Another
12 study available is a narrative review of global literature on these symptoms.⁴³ We have used
13 both subjective and objective assessment of loss of smell and taste to overcome the
14 shortcomings of using either method alone. Further, the individuals were blinded to the status
15 of COVID-19 at the time of examination, hence, their response to examination is not under
16 the influence of positive test result. Therefore, our estimates may be more closer to the actual
17 values. The solutions used for testing smell and taste were chosen to be culturally appropriate
18 for the population being tested and can be replicated using the same standardised
19 concentrations throughout the country.

20 **4.5. Limitations**

21 Our study has few limitations. The clinical examination of sense of smell and taste still
22 entails a subjective component by relying on the participant's responses. We attempted to
23 detect and eliminate this bias by using a control solution, and did not find any positive
24 reporting of smell or taste with control solution. We did not measure odour threshold, which

1 has been reported to be more affected in COVID-19 affected persons compared to odour
2 perception and discrimination. Since we did not include hospitalised patients being tested for
3 COVID-19, we are likely to have included only mild and moderate cases of COVID-19 in our
4 study. Since the number of COVID-19 affected person reporting these symptoms was small,
5 we could not comment on the duration of the symptoms or explore the association of these
6 symptoms with other risk factors, disease severity or with the outcomes of the infection.

7 **Financial support & Sponsorship:**

8 Indian Council of Medical Research- National Institute of Epidemiology (ICMR-NIE)

9

10 **Conflict of interest** - The authors have no conflicts of interest to disclose.

11 **Acknowledgement**

12 The study was funded by the Indian Council of Medical Research- National Institute of
13 Epidemiology, Chennai, India. This organization functions under the Department of Health
14 Research, Ministry of Health and Family Welfare. The authors (KJ, MR, PM, SM, AR, RRG,
15 MM) are affiliated to this institute. The corresponding author (MM) hereby states that he has
16 full access to all the data in the study, and had final responsibility for the decision to submit
17 for publication.

18 The authors also wish to acknowledge the contribution of Dr.S.Ramu, Neurologist,
19 Sivagangai Medical College, Tamil Nadu, India for his guidance in designing the
20 psychophysical assessment of olfaction and gustation.

21

1

2 **References**

3

- 4 1. Coronavirus disease (COVID-19) [Internet]. [cited 2020 Aug 26]. Available from:
5 [https://www.who.int/emergencies/diseases/novel-coronavirus-](https://www.who.int/emergencies/diseases/novel-coronavirus-2019?gclid=Cj0KCQjwvuj5BRDkARIsAGD9vIKMRuLJV17rS6z6QGgtKJZWnmKn9_Tzb8BYoDIR1R6ttIITmBfOkJ0aArGVEALw_wcB)
6 [2019?gclid=Cj0KCQjwvuj5BRDkARIsAGD9vIKMRuLJV17rS6z6QGgtKJZWnmKn9_Tzb8](https://www.who.int/emergencies/diseases/novel-coronavirus-2019?gclid=Cj0KCQjwvuj5BRDkARIsAGD9vIKMRuLJV17rS6z6QGgtKJZWnmKn9_Tzb8BYoDIR1R6ttIITmBfOkJ0aArGVEALw_wcB)
7 [BYoDIR1R6ttIITmBfOkJ0aArGVEALw_wcB](https://www.who.int/emergencies/diseases/novel-coronavirus-2019?gclid=Cj0KCQjwvuj5BRDkARIsAGD9vIKMRuLJV17rS6z6QGgtKJZWnmKn9_Tzb8BYoDIR1R6ttIITmBfOkJ0aArGVEALw_wcB)
- 8 2. Indian Council of Medical Research, New Delhi [Internet]. [cited 2020 Aug 14]. Available
9 from: <https://www.icmr.gov.in/>
- 10 3. Strategy of COVID19 testing in India. New Delhi: ICMR; 2020.
- 11 4. Revised Strategy of COVID19 testing in India (Version 3, dated 20/03/2020). New Delhi:
12 ICMR;
- 13 5. ADVISORY Newer Additional Strategies for COVID-19 Testing Existing strategies for
14 COVID-19 testing. New Delhi; 2020 Jun.
- 15 6. He J, Guo Y, Mao R, Zhang J. Proportion of asymptomatic coronavirus disease 2019
16 (COVID-19): a systematic review and meta-analysis. *J Med Virol*. 2020 Jul 21;
- 17 7. Adhikari SP, Meng S, Wu YJ, Mao YP, Ye RX, Wang QZ, et al. Epidemiology, causes,
18 clinical manifestation and diagnosis, prevention and control of coronavirus disease (COVID-
19 19) during the early outbreak period: A scoping review. Vol. 9, *Infectious Diseases of Poverty*.
20 BioMed Central Ltd.; 2020.
- 21 8. Guan W, Ni Z, Hu Y, Liang W, Ou C, He J, et al. Clinical Characteristics of Coronavirus
22 Disease 2019 in China. *N Engl J Med*. 2020 Feb 28;NEJMoa2002032.
- 23 9. Ge H, Wang X, Yuan X, Xiao G, Wang C, Deng T, et al. The epidemiology and clinical
24 information about COVID-19. *European Journal of Clinical Microbiology and Infectious*
25 *Diseases*. Springer; 2020. p. 1.
- 26 10. Fu L, Wang B, Yuan T, Chen X, Ao Y, Fitzpatrick T, et al. Clinical characteristics of
27 coronavirus disease 2019 (COVID-19) in China: A systematic review and meta-analysis. *J*
28 *Infect*. 2020 Jun 1;80(6):656–65.
- 29 11. Mao R, Qiu Y, He JS, Tan JY, Li XH, Liang J, et al. Manifestations and prognosis of
30 gastrointestinal and liver involvement in patients with COVID-19: a systematic review and
31 meta-analysis. *Lancet Gastroenterol Hepatol*. 2020;
- 32 12. Andina D, Noguera-Morel L, Bascuas-Arribas M, Gaitero-Tristán J, Alonso-Cadenas JA,
33 Escalada-Pellitero S, et al. Chilblains in children in the setting of COVID-19 pandemic.
34 *Pediatr Dermatol*. 2020 May 9;
- 35 13. Roland LT, Gurrola JG, Loftus PA, Cheung SW, Chang JL. Smell and taste symptom-based
36 predictive model for COVID-19 diagnosis. *Int forum allergy Rhinol*. 2020 May 4;
- 37 14. Symptoms of Coronavirus | CDC [Internet]. [cited 2020 May 22]. Available from:
38 <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html>

- 1 15. Xydakis MS, Dehgani-Mobaraki P, Holbrook EH, Geisthoff UW, Bauer C, Hautefort C, et al.
2 Smell and taste dysfunction in patients with COVID-19. *The Lancet Infectious Diseases*.
3 Lancet Publishing Group; 2020.
- 4 16. Gautier JF, Ravussin Y. A New Symptom of COVID-19: Loss of Taste and Smell. Vol. 28,
5 Obesity. Blackwell Publishing Inc.; 2020. p. 848.
- 6 17. Lechien JR, Chiesa-Estomba CM, De Siati DR, Horoi M, Le Bon SD, Rodriguez A, et al.
7 Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of
8 the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch*
9 *Otorhinolaryngol*. 2020 Aug 1;
- 10 18. Lechien JR, Chiesa-Estomba MD CM, Hans S, Barillari MD MR, Jouffe L, Saussez S. Loss of
11 Smell and Taste in 2013 European Patients With Mild to Moderate COVID-19. *Ann Intern*
12 *Med*. 2020 May 26;M20-2428.
- 13 19. Vaira LA, Salzano G, Fois AG, Piombino P, De Riu G. Potential pathogenesis of ageusia and
14 anosmia in COVID-19 patients. *International Forum of Allergy and Rhinology*. John Wiley
15 and Sons Inc.; 2020. p. alr.22593.
- 16 20. Vaira LA, Salzano G, Deiana G, De Riu G. Anosmia and Ageusia: Common Findings in
17 COVID-19 Patients. *Laryngoscope*. 2020;
- 18 21. Tong JY, Wong A, Zhu D, Fastenberg JH, Tham T. The Prevalence of Olfactory and
19 Gustatory Dysfunction in COVID-19 Patients: A Systematic Review and Meta-analysis.
20 *Otolaryngol Neck Surg*. 2020 May 5;019459982092647.
- 21 22. Le Bon SD, Pisarski N, Verbeke J, Prunier L, Cavelier G, Thill MP, et al. Psychophysical
22 evaluation of chemosensory functions 5 weeks after olfactory loss due to COVID-19: a
23 prospective cohort study on 72 patients. *Eur Arch Otorhinolaryngol*. 2020;
- 24 23. Whitcroft KL, Hummel T. Olfactory Dysfunction in COVID-19: Diagnosis and Management.
25 *JAMA - J Am Med Assoc*. 2020 Jun 23;323(24):2512-4.
- 26 24. Struyf T, Deeks JJ, Dinnes J, Takwoingi Y, Davenport C, Leeflang MMG, et al. Signs and
27 symptoms to determine if a patient presenting in primary care or hospital outpatient settings
28 has COVID-19 disease. Vol. 2020, *Cochrane Database of Systematic Reviews*. John Wiley
29 and Sons Ltd; 2020.
- 30 25. Menni C, Valdes AM, Freidin MB, Sudre CH, Nguyen LH, Drew DA, et al. Real-time
31 tracking of self-reported symptoms to predict potential COVID-19. *Nat Med*. 2020;
- 32 26. Vaira LA, Hopkins C, Petrocelli M, Lechien JR, Soma D, Giovanditto F, et al. Do olfactory
33 and gustatory psychophysical scores have prognostic value in COVID-19 patients? A
34 prospective study of 106 patients. *J Otolaryngol - head neck Surg = Le J d'oto-rhino-*
35 *laryngologie Chir cervico-faciale*. 2020 Aug 6;49(1):56.
- 36 27. Campbell, William Wesley DeJong RN. *DeJong's the Neurologic Examination*. Lippincott
37 Williams & Wilkins; 2012.
- 38 28. Abraham P, Aggarwal N, Babu GR, Barani S, Bhargava B, Bhatnagar T, et al. Laboratory
39 surveillance for SARS-CoV-2 in India: Performance of testing & descriptive epidemiology of
40 detected COVID-19, January 22-April 30, 2020. *Indian J Med Res*. 2020 May 1;151(5):424-
41 37.

- 1 29. Jutzeler CR, Bourguignon L, Weis C V., Tong B, Wong C, Rieck B, et al. Co-morbidities,
2 clinical signs and symptoms, laboratory findings, imaging features, treatment strategies, and
3 outcomes in adult and pediatric patients with COVID-19: A systematic review and meta-
4 analysis. *Travel Med Infect Dis.* 2020 Aug 4;101825.
- 5 30. Hoang A, Chorath K, Moreira A, Evans M, Burmeister-Morton F, Burmeister F, et al.
6 COVID-19 in 7780 pediatric patients: A systematic review. *EClinicalMedicine.* 2020 Jul 1;24.
- 7 31. Rodriguez-Morales AJ, Cardona-Ospina JA, Gutiérrez-Ocampo E, Villamizar-Peña R,
8 Holguin-Rivera Y, Escalera-Antezana JP, et al. Clinical, laboratory and imaging features of
9 COVID-19: A systematic review and meta-analysis. Vol. 34, *Travel Medicine and Infectious*
10 *Disease.* Elsevier USA; 2020.
- 11 32. Mullol J, Alobid I, Mariño-Sánchez F, Izquierdo-Domínguez A, Marin C, Klimek L, et al. The
12 Loss of Smell and Taste in the COVID-19 Outbreak: a Tale of Many Countries. Vol. 20,
13 *Current Allergy and Asthma Reports.* Springer; 2020.
- 14 33. Carrillo-Larco RM, Altez-Fernandez C. Anosmia and dysgeusia in COVID-19: A systematic
15 review. Vol. 5, *Wellcome Open Research.* F1000 Research Ltd; 2020.
- 16 34. Samaranayake LP, Fakhruddin KS, Panduwawala C. Sudden onset, acute loss of taste and
17 smell in coronavirus disease 2019 (COVID-19): a systematic review. *Acta Odontol Scand.*
18 2020 Aug 7;1-7.
- 19 35. Agyeman AA, Lee Chin K, Landersdorfer CB, Liew D, Ofori-Asenso R. Smell and Taste
20 Dysfunction in Patients With COVID-19: A Systematic Review and Meta-analysis. *Mayo Clin*
21 *Proc.* 2020 Jun;95(8).
- 22 36. Borsetto D, Hopkins C, Philips V, Obholzer R, Tirelli G, Polesel J, et al. Self-reported
23 alteration of sense of smell or taste in patients with COVID-19: a systematic review and meta-
24 analysis on 3563 patients. *Rhinology.* 2020;1-7.
- 25 37. Boscolo-Rizzo P, Borsetto D, Spinato G, Fabbris C, Menegaldo A, Gaudio P, et al. New
26 onset of loss of smell or taste in household contacts of home-isolated SARS-CoV-2-positive
27 subjects. *Eur Arch Otorhinolaryngol.* 2020;
- 28 38. Rocke J, Hopkins C, Philpott C, Kumar N. Is loss of sense of smell a diagnostic marker in
29 COVID-19: A Systematic Review and Meta-analysis. *Clin Otolaryngol.* 2020
30 Aug;coa.13620.
- 31 39. Kucirka LM, Lauer SA, Laeyendecker O, Boon D, Lessler J. Variation in False-Negative Rate
32 of Reverse Transcriptase Polymerase Chain Reaction-Based SARS-CoV-2 Tests by Time
33 Since Exposure. *Ann Intern Med.* 2020 Aug 18;173(4):262-267. doi: 10.7326/M20-1495.
34 Epub 2020 May 13. PMID: 32422057; PMCID: PMC7240870.
- 35 40. Desiato VM, Levy DA, Byun YJ, Nguyen SA, Soler ZM, Schlosser RJ. The Prevalence of
36 Olfactory Dysfunction in the General Population: A Systematic Review and Meta-analysis.
37 *Am J Rhinol Allergy.* 2020 Aug 3;194589242094625.
- 38 41. Lechien JR, Cabaraux P, Chiesa-Estomba CM, Khalife M, Hans S, Calvo-Henriquez C, et al.
39 Objective olfactory evaluation of self-reported loss of smell in a case series of 86 COVID-19
40 patients. In: *Head and Neck.* John Wiley and Sons Inc.; 2020.
- 41 42. Hannum ME, Ramirez VA, Lipson SJ, Herriman RD, Toskala AK, Lin C, et al. Objective
42 sensory testing methods reveal a higher prevalence of olfactory loss in COVID-19 positive

1 patients compared to subjective methods: A systematic review and meta-analysis. medRxiv
2 Prepr Serv Heal Sci. 2020 Jul 6;

3 43. Harikrishnan P. Gustatory Dysfunction as an Early Symptom in COVID-19 Screening. J
4 Craniofac Surg. 2020 Jul 7;

5

6

7

8

9

10

11

12

13 Box 1. Standardisation of testing kits and procedures for testing of sense of smell and
14 taste

15 Serially increasing concentrations of the solutions to be used for smell and taste testing were
16 prepared at the laboratory. The solutions were presented to a group of 10 apparently healthy
17 volunteers. They were required to indicate the concentration at which they perceived the
18 particular smell or taste. A supra-threshold concentration i.e., a concentration higher than the
19 highest concentration at which the smell or taste was perceived by all volunteers was chosen
20 for the study participants. The concentration used in the study were as follows:

Substance	Concentration
For gustation	
Sugar	32g/100ml
Lemon concentrate	20ml/100 ml
Salt	2.5g/100 ml
For olfaction	
Coffee	6g/ 100ml
Asafoetida	2.5g/100 ml
Eucalyptus oil	5 ml concentrate undiluted

21

1 The order of odour presented to the participants was Asafoetida, followed by Coffee and
2 Eucalyptus oil in both the nostrils, one after the other. Drinking water was used as control
3 solution. Cotton buds dipped in the respective solutions were kept close to the opening of the
4 nostril and the participant was asked to identify the odour. Separate buds were used for all the
5 patients and disposed appropriately after single use. Nearly 10-15 seconds was given for
6 desensitization between different odours.

7 We used salt, sugar and lemon extract solutions to identify the bitter, sweet and sour tastes,
8 respectively. The solutions were prepared freshly on the morning of each day and kept for 3
9 hours before preparing again. Drinking water was used as control solution. We used single-
10 use lolly sticks to dip into the solutions and apply on the participant's tongue for 10-15
11 seconds before asking them to identify the correct taste. Drinking water was given before and
12 after each solution to rinse the mouth.

13 The data collectors followed the Ministry of Health and Family Welfare, India guidelines for
14 donning and doffing complete personal protective equipment.

15

16

Table 1: Baseline characteristics persons tested for COVID -19, Chennai, Tamil Nadu

Characteristics		Total (N=277)		COVID-19 Negative (N=219)		COVID-19 Positive (N=58)	
		n	%	n	%	n	%
Gender	Male	169	61.0	139	63.5	30	51.7
	Female	108	39.0	80	36.5	28	48.3
Read and write	Yes	247	93.2	195	93.3	52	92.9
	No	18	6.8	14	6.7	4	7.1
Occupation exposure to strong fumes*	Yes	9	3.4	6	2.9	3	5.4
	No	255	96.2	203	97.1	52	92.9
Income	Below Poverty Line*	20	7.5	15	7.2	5	8.9
	Above Poverty Line	213	80.4	172	82.3	41	73.2
	No response	32	12.1	22	10.5	10	17.9
Patient	Symptomatic contact	16	5.8%	11	5.0%	5	8.6%

category	Symptomatic Healthcare worker / Frontline workers	8	2.9%	8	3.7%	0	0.0%
	Asymptomatic family member	94	33.9%	80	36.5%	14	24.1%
	Asymptomatic healthcare worker	26	9.4%	25	11.4%	1	1.7%
	Symptomatic Influenza like Illness (ILI)	21	7.6%	12	5.5%	9	15.5%
	Pregnant woman in / near labour	1	0.4%	0	0.0%	1	1.7%
	Symptomatic among returnees and migrants	32	11.6%	23	10.5%	9	15.5%
	Persons from Hotspot / Containment zones.	79	28.5%	60	27.4%	19	32.8%

*A monthly household income of less than 6000 was operationally defined as below poverty line, data was missing for 12 persons on

occupational exposure, income and education

Table 2. Behavioral risk factors and comorbidities among persons tested for COVID - 19, Chennai, Tamil Nadu

Characteristics	Total (N=277)		COVID-19 Negative (N=219)		COVID-19 Positive (N=58)	
	n	%	n	%	n	%
Smokeless tobacco	13	4.9	12	5.7	1	1.8
Smoking	27	10.2	21	10.0	6	10.7
Alcohol	32	12.1	28	13.4	4	7.1
Hypertension	27	10.2%	17	8.1%	10	17.9%
Diabetes mellitus	30	11.3%	20	9.6%	10	17.9%
Heart diseases	6	2.3%	6	2.9%	0	0.0%
Asthma	4	1.5%	4	1.9%	0	0.0%
Chronic kidney diseases	0	0.0%	0	0.0%	0	0.0%
Liver diseases	3	1.1%	3	1.4%	0	0.0%

Neurological disorder	3	1.1%	2	1.0%	1	1.8%
Rheumatologic disorder	1	0.4%	1	0.5%	0	0.0%
ENT structural abnormalities	5	1.9%	3	1.4%	2	3.6%
Other comorbidities	15	5.7%	11	5.3%	4	7.1%

Table 3. Prevalence (95% Confidence Interval) of loss of smell, loss of taste among patients tested for COVID -19, Chennai, Tamil Nadu

Parameter	Method of examination	Total (N=277)		COVID-19 Negative (N=219)		COVID-19 Positive (N=58)	
		n (%)	95% CI	n (%)	95% CI	n (%)	95 %CI
Loss of smell	SR	19 (6.9%)	4.2-10.5	14 (6.4%)	3.5-10.5	5 (8.6%)	2.9-19.0
	CE	15 (5.4%)	3.1-8.8	9 (4.1%)	1.9-7.7	6 (10.3%)	3.9-21.2
	SR / CE	31 (11.2%)	8.3-16.5	21 (9.6%)	6.5 -13.2	10 (17.2)	9.1-30.1
Loss of taste	SR	15 (5.4%)	3.1-8.8	12 (5.5%)	2.8-9.4	3 (5.2%)	1.1-14.4
	CE	9 (3.2%)	1.5-6.1	8 (3.7%)	1.5-7.1	1 (1.7%)	0.0-9.2
	SR / CE	24 (8.7%)	6.1-13.6	20 (9.1%)	6.1-14.8	4 (6.9%)	2.1-17.9
Loss of smell or taste	SR	21 (7.6%)	5.1-12.1	14 (6.4%)	3.8-11.2	7 (12.1%)	5.3-27.5
	CE	18 (6.5%)	4.3-11.0	11 (5.0)	2.8-9.6	7 (12.1%)	5.5-25.4
	SR / CE	36 (13.0%)	9.9-18.7	23 (10.5)	7.3-16.4	13 (22.4%)	13.2-31.0

*SR-self reported; CE- Clinical examination

Table 4. Positive and Negative predictive values of the symptoms loss of smell and taste for COVID 19 status among patients tested for COVID -19, Chennai, Tamil Nadu

Parameter	Method of examination*	NPV	PPV	Likelihood Ratio of a positive test
Loss of smell	SR	79.5%	26.3%	1.34
	CE	80.2%	40.0%	2.51
	SR / CE	80.5%	32.3%	1.79
Loss of taste	SR	79.0%	20.0%	0.95
	CE	78.7%	11.1%	0.46
	SR / CE	78.7%	16.7%	0.76
Loss of smell or taste	SR	80.1%	33.3%	1.89
	CE	80.3%	38.9%	2.42
	SR / CE	81.3%	18.7%	2.13

*SR-self reported; CE- Clinical examination

Figure 1: Proportion reporting different symptoms among patients tested for COVID - 19 at Chennai, Tamil Nadu

