



408/D

Pathogenic *Leptospira* spp induce higher superoxide anion production by HL 60-derived neutrophils

T R G N Fernando,¹ M J R Niloofa,¹ H J de Silva,² S Rajapakse,³ S Premawansa⁴ and S M Handunnetti¹

¹*Institute of Biochemistry, Molecular Biology and Biotechnology, University of Colombo, Colombo 03,*

²*Faculty of Medicine, University of Kelaniya, Kelaniya*

³*Tropical Medicine Research Unit, Faculty of Medicine, University of Colombo, Colombo 08,*

⁴*Faculty of Science, University of Colombo, Colombo 03*

The superoxide anion (O_2^-) is the first free radical produced in phagocytes which enables pathogen killing and also generates other microbicidal reactive oxygen species (ROS). The induction of high O_2^- production by phagocytes as a virulence factor for pathogenic *Leptospira* is not known. The objective of this study was to assess the induced O_2^- production by human neutrophils exposed *Leptospira* and compare the differences in the level of O_2^- produced by HL-60 derived neutrophils interacted with pathogenic and saprophytic *Leptospira* spp. HL-60 cells were treated with dimethyl sulfoxide to differentiate in to mature neutrophils. Saprophytic *Leptospira biflexa* serovar Patoc and pathogenic *L. Interrogans* serovar Pyrogenes were grown in Ellinghausen-McCullough-Johnson-Harris media containing 10% heat inactivated rabbit sera. Intracellular O_2^- production of neutrophils interacted with bacteria in the presence of sera from leptospirosis patients (n=30/group of severe and mild leptospirosis patients) and healthy subjects (n=20) was assessed using quantitative nitroblue tetrazolium assay. Both anti-leptospiral IgM and IgG levels were measured in test sera prior to the assay. In the presence of normal and also patients' sera, neutrophils interacted with *L. Interrogans* Pyrogenes produced significantly high O_2^- levels than that of *L. biflexa* (p<0.05). When *Leptospira* density was varied for neutrophil-bacteria interactions, low *Leptospira* densities induced higher O_2^- productions compared to the high concentrations (p<0.03). Moreover, analysis of O_2^- production by neutrophils in the presence of patients' sera showed that, opsonization of both saprophytic and pathogenic *Leptospira* by IgM or IgG had not induced a significant difference in the level of O_2^- produced (p = 0.012). These results suggest that, the increased O_2^- production induced by *L. Interrogans* Pyrogenes could be a virulent factor in pathogenic *Leptospira* spp which may contribute to the pathogenesis of severe leptospirosis.

Acknowledgement: National Research Council- NRC-12-077